



**Table H. Annual Incidence Rates of Selected Notifiable Diseases by Year of Onset  
Los Angeles County, 2011-2016**

Disease	Annual Incidence Rate (Cases per 100,000) <sup>b</sup>					
	2011	2012	2013	2014	2015	2016
Amebiasis	0.93	1.06	0.61	0.68	0.65	0.73
Botulism	0.03	0.04	0.04	0.01	0.02	0.06
Brucellosis	0.06	0.04	0.11	0.07	0.08	0.06
Campylobacteriosis	13.60	16.63	18.11	15.93	16.96	16.29
Cholera	-	-	-	-	0.04	-
Coccidioidomycosis	3.28	3.52	3.85	4.51	6.40	8.43
Cryptosporidiosis	0.55	0.47	0.51	0.83	0.59	1.02
Cysticercosis	0.40	0.12	0.01	0.10	0.13	0.06
Dengue	-	0.02	0.02	0.34	0.31	0.48
<i>E. Coli</i> —Shiga Toxin-Producing	0.95	1.04	1.08	0.95	1.83	2.94
Encephalitis	0.64	0.81	0.84	0.97	1.42	0.72
Giardiasis	3.15	3.16	4.17	3.66	3.96	4.71
Hansen’s Disease (Leprosy)	0.02	0.03	0.01	0.03	-	0.01
Hepatitis A	0.49	0.51	0.64	0.44	0.34	0.69
Hepatitis B	0.65	0.41	0.58	0.44	0.52	0.44
Hepatitis C	0.11	0.08	0.05	0.05	0.02	0.05
Hepatitis Unspecified	0.04	-	-	-	-	-
Legionellosis	1.25	1.19	0.90	1.48	1.79	2.55
Listeriosis, Nonperinatal	0.21	0.28	0.24	0.29	0.36	0.34
Listeriosis, Perinatal <sup>a</sup>	4.95	5.71	3.34	4.11	2.58	3.48
Lyme Disease	0.06	0.01	0.12	0.05	0.04	0.01
Malaria	0.24	0.20	0.17	0.22	0.28	0.25
Meningitis, Viral	3.42	3.26	3.77	4.23	3.83	1.91
Meningococcal Infections	0.40	0.13	0.18	0.12	0.13	0.21
Pneumococcal Disease, Invasive	7.11	5.42	5.55	4.87	4.89	5.24
Psittacosis	-	-	-	-	-	-
Q-fever	-	0.03	0.02	0.01	0.05	0.02
Relapsing Fever	-	-	-	0.01	-	-
Rheumatic Fever, Acute	-	-	-	-	-	-
Salmonellosis	9.72	11.20	10.74	12.07	11.95	10.91
Shigellosis	2.85	3.29	2.41	3.70	5.31	6.08
Streptococcus, Group A Invasive	1.89	1.81	2.07	2.35	2.37	3.68
Strongyloidiasis	-	-	0.12	0.37	0.09	0.10
Taeniasis	0.05	0.06	0.04	0.03	0.02	0.02
Tetanus	-	-	0.01	-	-	-
Trichinosis	-	-	-	-	-	-
Tularemia	-	-	-	-	-	-
Typhoid Fever, Case	0.16	0.06	0.18	0.16	0.15	0.11
Typhoid Fever, Carrier	0.03	-	-	-	-	0.02
Typhus Fever	0.41	0.54	0.72	0.47	0.56	0.49
Vibrio	0.21	0.31	0.28	0.55	0.45	0.34
West Nile Virus	0.68	1.87	1.75	2.31	3.13	1.59

<sup>a</sup>Rates for perinatal listeriosis were calculated as cases per 100,000 live births.

<sup>b</sup>Rates of disease based on less than 19 cases or events are considered "unreliable." A zero rate made from no events is especially hazardous and are not reported here, except with a dash ("-"). Conclusions drawn from unreliable rates should be made with caution, if they are to be made at all.