


# SHIGA-TOXIN PRODUCING *ESCHERICHIA COLI* STEC Update



Roshan Reporter, MD, MPH

Rita Bagby, PS-PHN

---

Leticia Martinez, PS-PHN



# Objectives


**At the conclusion of this presentation the participant should be able to:**

- Know the clinical signs and symptoms of the disease caused by Shiga toxin-producing *Escherichia coli* (STEC)
  - Know the epidemiology of STEC infection in Los Angeles County.
  - Know the case management of patients with STEC infection.
  - Identify laboratory tests used to diagnose Shiga toxin-producing *Escherichia coli* infections
- 



# WHAT IS *ESCHERICHIA COLI*?

- Gram Negative Bacteria
- Sources can be: Urine, Resp, Blood, and Stool
- Considered normal flora in intestines of many mammals including humans
- **Some** E. coli causes GI disease
  - These are pathogenic E.coli; they possess ability to produce toxin
  - STEC is the type of E.coli are the topic today.



# What are Shiga Toxin-producing E.coli ?

- Certain bacteria produce a toxin called shiga toxin: some E.coli can do this
- These E. coli are called “Shiga toxin-producing” E. coli, or STEC.
- You may hear them called verocytotoxic E.coli (VTEC) or enterohemorrhagic E.coli (EHEC)



# INCUBATION

The incubation period is usually 3-4 days after the exposure, but may be as short as 1 day or as long as 10 days



# Signs and Symptoms

- Diarrhea (blood is common)
- Abdominal cramps (usually severe)
- Little or no Fever (less than 101F)




# Pathophysiology

- Infection
  - Body's response
  - Organism is replicating and producing toxin
- Effects of toxin(s)
  - Adheres to cell
    - Epithelial intestinal
    - Endothelial Blood vessels
    - Renal Endothelial cells
    - Red blood cells



# Sequence of events in STEC infection





# Major modes of Transmission

- Food
  - cattle products, e.g., beef, raw milk
  - food contaminated with cattle or human feces e.g., lettuce, spinach, and cookie dough.
- Water
  - Drinking water
  - Recreational water
- Animal contact
  - contact with farm animals, e.g. petting zoos
  - contact with farm animals' environment
- Human contact
  - With the feces of infected persons



# Transmission




Modes of transmission in  
non-O157 STEC outbreaks,  
U.S., 1990-2007  
(N = 23)

<b>Mode</b>	<b>No. outbreaks</b>
<b>Food</b>	<b>11</b>
<b>Person-to-person</b>	<b>6</b>
<b>Lake water</b>	<b>3</b>
<b>Animal contact</b>	<b>2</b>
<b>Undetermined</b>	<b>1</b>



# Seasonality of human non-O157 STEC isolates submitted to CDC, 1983-2002

(N=940 isolates)



Brooks, JID 2005



# Public Health Implications

- As infectious as *Shigella*
  - Low infectious dose (10-100)
  - SOS assessment critical
  - If congregate setting, assess for other ills
- Increased morbidity/mortality
  - Hemolytic Uremic Syndrome (HUS)
  - Thrombotic thrombocytopenic purpura (TTP)
- Outbreaks
  - Local
  - National





VS



STEAK

Ground Beef



# Special considerations

- Hamburger vs steak
- Pre washed vs washed
- Aged cheese vs fresh/soft cheese
- Pasteurized vs unpasteurized




# PHN Observations/Considerations

- Food preferences
- Food at home
- Kitchen technique
- Animal exposure
- SOS
- Patient Education/comprehension
- Remember the focus should be on the case and identifying any potential source.





# Two Possible Means of Transmission



# Prevention

- Wash your hands
  - Cook meats thoroughly.
  - Avoid unpasteurized milk/milk products
  - Avoid swallowing un-chlorinated water
  - Be cautious around animals
  - Wash produce
  - Avoid cross-contamination
- 




# FBI or No FBI?

- Focus on the incubation period
  - Ground Beef: Well cooked vs uncooked
  - Vegetarian
  - Kitchen technique
  - Food Preferences
  - FBI should be based on your best assessment of the situation
- 



# Pyramid of Surveillance

Reported to health department & CDC



# Milestones in STEC Follow-up

- 1994 *E. coli* O157 infection made reportable
  - 1995 Commercial Shiga toxin enzyme immunoassay (EIA) introduced
  - 2000 Non-O157 STEC infections made nationally reportable
  - 2009 CDC recommending that all diarrheal stool should be cultured for STEC and tested for the detection of Shiga toxins.
- 



# Testing for STEC using the Shiga toxin EIA

- Clinical lab processes stool specimen in broth
  - Tests broth for Shiga toxin using EIA
  - Positive test is reportable
- Clinical lab should send Shiga toxin-positive broth to Public Health lab
  - PH lab isolates STEC
  - PHL serotypes
  - If unable to serotype, will refer (State or CDC)



# Challenges related to use of the Shiga toxin EIA

- After adopting the EIA, some clinical labs stopped testing for *E. coli* O157 using selective media
    - *E. coli* O157 outbreaks could be missed
  - Some clinical labs discard Shiga toxin-positive specimens without obtaining an isolate, so
    - simply report “Shiga toxin positive” to doctor
    - serogroup not determined
- 
- *E. coli* O157 strains not identified and sub-typed for outbreak detection
  - Non-O157 outbreaks less likely identified



# *E. coli* serotyping


Lipopolysaccharide (LPS)

= O antigen

O1-O181

Flagella = H antigen

H1-H56



Slide from USDA presentaion





# In House PHL Testing

- O157
- O111
- O103
- O121
- O26



# Top Non-O157 Serotypes (CDC)

- O26            22% of non-O157 STEC
- O111          16% of non-O157 STEC
- O103          12% of non-O157 STEC
- O121          9% of non-O157 STEC
- O45            7% of non-O157 STEC
- O145          5% of non-O157 STEC



# Case History Form

- Why was it changed?
- Why is it better than the old form?
- What is new?
- What stayed the same ?



