



**Table G. Reported Cases of Selected Notifiable Diseases
by Year of Onset Los Angeles County, 1991-1996**

Disease	Year of Onset						Previous 5-Year Average	5-year 95% Upper Limit ^a
	1991	1992	1993	1994	1995	1996		
AIDS ^b	2,849	2,970	2,881	2,901	2824	2536	2,885	2,995
Amebiasis.	294	239	290	230	187	210	248	336
Botulism	0	0	1	3	2	4	1	4
Brucellosis	6	6	4	11	3	12	6	12
Campylobacteriosis ^c	1,318	1,396	1,385	1,301	1,401	1,731	1,360	1,452
Chlamydia	22,437	21,679	20,521	21,258	18,659	20,191	20,911	23,728
Cholera	2	3	3	4	2	0	3	4
Coccidioidomycosis,	43	98	106	98	76	68	84	134
Cryptosporidiosis	N/A	N/A	102	231	211	148	N/A	N/A
Cysticercosis	52	43	59	41	26	30	44	69
Dengue	1	0	0	0	2	0	1	2
<i>E. coli</i> 0157:H7	N/A	N/A	9	13	9	18	N/A	N/A
Encephalitis	35	35	59	36	59	33	45	71
Foodborne outbreaks	7	1	7	12	15	12	8	19
Giardiasis	1,600	1,622	1,616	1,105	940	951	1,377	2,021
Gonorrhea	14,767	14,001	11,080	8,802	7,807	5,723	11,291	17,317
<i>Haemophilus influenzae type b</i>	115	38	22	20	6	4	40	125
Hansen's Disease (Leprosy)	32	17	24	25	10	9	22	38
Hepatitis, A	1,221	1,196	1,040	1,193	1,062	1,301	1,142	1,308
Hepatitis, B	523	1,077	288	289	231	230	482	1,170
Hepatitis non-A non-B	103	255	148	115	205	231	165	290
Hepatitis unspecified	60	45	32	40	22	28	40	68
Kawasaki syndrome	61	47	49	46	39	20	48	64
Legionellosis	N/A	16	11	11	16	12	N/A	N/A
Listeriosis nonperinatal	38	23	25	26	26	30	28	39
Listeriosis perinatal	24	22	23	11	10	5	18	32
Lyme disease	3	5	0	2	5	2	3	7
Malaria	51	69	40	40	76	59	55	88
Measles.	800	37	51	16	7	2	182	860
Meningitis viral	225	936	490	250	167	185	414	1,035
Meningococcal infection	115	85	98	91	52	59	88	134
Mumps	70	57	48	51	42	37	54	74
Pertussis	43	140	104	58	103	120	90	166
Psittacosis	0	1	1	0	0	0	0	1
Q-fever	1	0	0	0	0	0	0	1
Relapsing fever	0	1	1	0	0	0	0	1
Rheumatic fever, acute	1	3	3	5	1	2	3	6
Rubella.	53	6	3	4	3	2	14	57
Salmonellosis	1,582	1,617	1,565	2,091	2,084	1,757	1,788	2,325
Shigellosis	1,497	1,882	1,532	1,336	1,747	1,128	1,599	2,022
Strongyloidiasis	45	45	42	24	17	11	35	60
Syphilis. (early latent)	2,632	2,050	1,660	1,024	926	744	1,658	3,058
Syphilis (prim. & secon.)	1,256	726	511	326	271	216	618	1,399
Syphilis, (congenital)	448	322	257	202	161	126	278	499
Tetanus	0	1	1	0	4	1	1	4



Chapter from the *Communicable Disease Morbidity Report 1996*, Disease Control Programs.
County of Los Angeles Department of Health Services.

Trichinosis	0	2	0	2	1	0	1	3
Tuberculosis	2,121	2,198	1,940	1,794	1,622	1,375	1,935	2,396
Tularemia	0	1	0	0	1	1	0	1
Typhoid fever, case	48	45	42	38	26	31	40	57
Typhoid fever, carrier	15	8	8	5	6	4	8	16
Typhus fever,	4	0	14	8	11	16	7	18
Vibrio.	N/A	N/A	14	13	12	23	N/A	N/A

^aThe normal distribution assumption may not apply to some rare diseases.

^bAll data are adjusted for report delay and the 1993 AIDS case definition change.

^c1996 data over 95% upper limit.



**Table H. Annual Incidence Rates of Selected Notifiable Diseases
by Year of Onset Los Angeles County, 1991-1996**

Disease	Annual Incidence Rate (Cases per 100,000)					
	1991	1992	1993	1994	1995	1996
AIDS ^a	35.04	34.52	32.76	33.51	32.26	28.56
Amebiasis	3.62	2.78	3.30	2.66	2.14	2.36
Botulism	0.00	0.00	0.01	0.03	0.02	0.05
Brucellosis	0.07	0.07	0.05	0.13	0.03	0.14
Campylobacteriosis	16.21	16.23	15.75	15.03	16.00	19.49
Chlamydia	275.93	252.01	233.31	245.57	213.15	227.40
Cholera	0.02	0.03	0.03	0.05	0.02	0.00
Coccidioidomycosis	0.53	1.14	1.21	1.13	0.87	0.77
Cryptosporidiosis	N/A	N/A	1.16	2.67	2.41	1.67
Cysticercosis	0.64	0.50	0.67	0.47	0.30	0.34
Dengue	0.01	0.00	0.00	0.00	0.02	0.00
<i>E coli</i> O157:H7	N/A	N/A	0.10	0.15	0.10	0.20
Encephalitis	0.43	0.41	0.67	0.42	0.67	0.37
Foodborne outbreaks	0.09	0.01	0.08	0.14	0.17	0.14
Giardiasis	19.68	18.86	18.37	12.76	10.74	10.71
Gonorrhea	181.61	162.76	125.97	101.68	89.18	64.50
<i>Haemophilus influenzae</i> type b	1.41	0.44	0.25	0.23	0.07	0.05
Hansen's Disease (Leprosy)	0.39	0.20	0.27	0.29	0.11	0.10
Hepatitis A	15.02	13.90	11.82	13.78	12.13	14.65
Hepatitis B	6.43	12.52	3.27	3.34	2.64	2.59
Hepatitis non-A non-B	1.27	2.96	1.68	1.33	2.34	2.60
Hepatitis unspecified	0.74	0.52	0.36	0.46	0.25	0.32
Kawasaki syndrome	0.75	0.55	0.56	0.53	0.45	0.23
Legionellosis	N/A	0.19	0.13	0.13	0.18	0.14
Listeriosis, nonperinatal	0.47	0.27	0.28	0.30	0.30	0.34
Listeriosis, perinatal ^b	12.72	11.98	13.03	6.01	5.81	3.19
Lyme Disease	0.04	0.06	0.00	0.02	0.06	0.02
Malaria	0.63	0.80	0.45	0.46	0.87	0.66
Measles	9.84	0.43	0.58	0.18	0.08	0.02
Meningitis, viral	2.77	10.88	5.57	2.89	1.91	2.08
Meningococcal infections	1.41	0.99	1.11	1.05	0.59	0.66
Mumps	0.86	0.66	0.55	0.59	0.48	0.42
Pertussis	0.53	1.63	1.18	0.67	1.18	1.35
Psittacosis	0.00	0.01	0.01	0.00	0.00	0.00
Q-fever	0.01	0.00	0.00	0.00	0.00	0.00
Relapsing fever	0.00	0.01	0.01	0.00	0.00	0.00
Rheumatic fever, acute	0.01	0.03	0.03	0.06	0.01	0.02
Rubella	0.65	0.07	0.03	0.05	0.03	0.02
Salmonellosis	19.46	18.80	17.79	24.16	23.81	19.79
Shigellosis	18.41	21.88	17.42	15.43	19.96	12.70
Strongyloidiasis	0.55	0.52	0.48	0.28	0.19	0.12
Syphilis (early latent)	32.37	23.83	18.87	11.83	10.58	8.40
Syphilis (prim. & secon.)	15.45	8.44	5.81	3.77	3.10	2.40
Syphilis (congenital) ^c	237.51	175.47	145.70	125.60	98.10	76.80
Tetanus	0.00	0.01	0.01	0.00	0.05	0.01
Trichinosis	0.00	0.02	0.00	0.02	0.01	0.00
Tuberculosis	26.08	25.55	22.06	20.72	18.53	15.48



Chapter from the *Communicable Disease Morbidity Report 1996*, Disease Control Programs.
County of Los Angeles Department of Health Services.

Tularemia	0.00	0.01	0.00	0.00	0.01	0.01
Typhoid fever, case	0.59	0.52	0.48	0.44	0.30	0.35
Typhoid fever, carrier	0.18	0.09	0.09	0.06	0.07	0.05
Typhus fever	0.05	0.00	0.16	0.09	0.13	0.18
<i>Vibrio</i>	N/A	N/A	0.16	0.15	0.14	0.26

^aAll data are adjusted for report delay and 1993 AIDS case definition change

^bRates for perinatal Listeriosis and congenital syphilis were calculated as cases per 100,000 live births.



**Table I. Five-year Average of Notifiable Diseases by Month of Onset
Los Angeles County, 1992-1996**

Disease	Jan	Feb	Mar	Apr	May	Jun	July	Aug	Sept	Oct	Nov	Dec	Total^a
AIDS	59.2	51.2	60.8	53.6	52.4	53.0	50.0	50.4	49.2	51.2	42.6	46.2	619.8
Amebiasis	18.2	15.6	20.2	20.4	18.2	18.0	22.2	20.6	19.0	17.6	13.6	17.0	220.6
Botulism	0.2	0.2	0.0	0.0	0.4	0.0	0.0	0.4	0.2	0.2	0.0	0.2	1.8
Brucellosis	0.6	0.2	0.4	0.6	0.4	0.8	1.4	0.4	0.6	0.8	0.2	0.2	6.6
Campylobacteriosis	78.8	74.0	82.8	116.4	148.2	153.4	163.0	137.2	130.8	113.6	105.6	84.4	1388.2
Cholera	0.2	0.2	0.2	0.4	0.4	0.2	0.2	0.0	0.0	0.0	0.0	0.0	1.8
Coccidioidomycosis	6.2	9.0	6.2	6.4	4.8	3.8	6.6	6.2	7.2	8.2	6.4	7.2	78.2
Cryptosporidiosis ^b	13.3	11.5	12.5	8.0	12.3	11.3	14.0	12.8	22.5	19.8	13.5	15.3	166.5
Cysticercosis	3.8	2.8	4.2	2.2	2.8	3.6	3.4	2.2	3.4	2.2	3.8	1.6	36.0
Dengue	0.2	0.0	0.0	0.0	0.0	0.2	0.0	0.0	0.0	0.0	0.0	0.0	0.4
<i>E. coli</i> 0157:H7 ^b	0.3	0.8	0.5	1.0	1.0	2.3	0.3	1.0	1.0	2.5	1.5	0.5	12.5
Encephalitis	2.6	2.4	3.6	3.6	4.2	4.2	2.6	2.6	2.2	1.6	3.2	3.0	35.8
Foodborne outbreaks	0.4	0.6	0.8	0.6	1.0	1.2	0.6	1.2	1.0	0.8	0.4	0.8	9.4
Giardiasis	72.0	64.4	95.6	85.4	96.2	98.6	115.2	138.6	129.6	119.2	83.0	68.8	1166.6
<i>Haemophilus influenzae</i> type b	4.2	3.6	3.2	2.4	2.0	1.8	1.2	1.4	2.2	2.6	2.0	4.2	31.0
Hansen's Disease(Leprosy)	3.0	1.0	2.4	1.2	0.8	2.2	1.8	0.4	1.0	0.0	0.8	0.6	15.2
Hepatitis A	75.8	70.8	73.6	63.6	73.0	78.8	81.4	89.2	115.2	118.6	109.0	91.2	1040.2
Hepatitis B	25.4	21.4	27.8	31.0	29.2	32.6	30.2	29.8	30.6	31.2	23.8	23.4	336.4
Hepatitis non-A non-B	13.6	11.8	14.4	17.0	13.8	15.2	14.8	14.6	11.8	16.2	12.8	16.8	172.8
Hepatitis unspecified	2.4	1.8	3.4	2.2	1.8	3.0	3.0	1.2	2.6	3.0	1.2	1.2	26.8
Kawasaki syndrome	4.6	5.6	4.8	4.6	3.2	1.2	2.4	1.8	2.6	2.8	2.8	2.6	39.0
Legionellosis	0.4	0.8	2.8	1.2	0.8	0.4	1.6	1.4	1.0	0.4	0.4	1.0	12.0
Listeriosis, nonperinatal	1.2	1.8	1.8	1.6	2.2	2.8	2.0	4.4	2.2	2.8	1.4	1.4	25.6
Listeriosis, perinatal	1.0	1.2	1.2	1.4	1.2	1.6	1.2	0.8	2.2	1.6	0.4	0.2	14.0
Lyme Disease	0.0	0.0	0.2	0.0	0.6	0.0	0.6	0.2	0.2	0.2	0.0	0.0	2.0
Malaria	3.4	3.0	2.2	4.0	7.2	3.6	4.8	6.4	6.6	3.8	3.0	3.6	51.6
Measles	1.8	1.2	0.8	2.0	5.0	5.2	4.2	1.6	1.0	0.6	0.0	0.4	23.8
Meningitis, viral	12.8	11.2	16.0	20.6	29.8	43.4	55.2	78.2	51.0	42.4	23.8	11.0	395.4
Meningococcal infections	8.4	10.4	6.4	4.6	4.6	5.8	3.2	4.4	2.4	2.4	2.6	7.0	62.2
Mumps	4.2	4.2	4.4	4.4	6.6	3.0	2.4	3.2	4.4	3.8	2.0	1.0	43.6
Pertussis	6.0	4.8	6.2	7.0	8.4	9.6	11.2	13.4	11.6	11.0	9.0	6.8	105.0
Psittacosis	0.0	0.0	0.0	0.2	0.0	0.2	0.0	0.0	0.0	0.0	0.0	0.0	0.4
Relapsing fever	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.2	0.2	0.0	0.0	0.4
Rheumatic fever, acute	0.4	0.2	0.0	0.8	0.0	0.0	0.0	0.2	0.4	0.4	0.0	0.0	2.4
Rubella	0.0	0.0	1.0	0.2	0.4	0.0	0.6	0.2	0.4	0.8	0.2	0.0	3.8
Salmonellosis	83.6	81.4	124.0	128.4	153.8	155.6	187.4	236.2	208.0	177.6	132.2	97.6	1765.8
Shigellosis	70.0	56.8	68.2	77.4	94.6	128.0	190.0	248.8	222.0	153.8	92.8	69.0	1471.4
Strongyloidiasis	1.6	1.2	1.2	2.8	2.4	3.4	2.0	2.4	2.8	2.0	2.4	1.0	25.2
Tetanus	0.2	0.2	0.0	0.2	0.4	0.0	0.0	0.2	0.0	0.0	0.0	0.2	1.4
Tularemia	0.0	0.0	0.0	0.0	0.0	0.0	0.2	0.2	0.0	0.2	0.0	0.0	0.6
Typhoid fever, case	2.4	3.4	2.8	2.4	5.2	4.0	3.8	5.2	3.0	1.8	1.4	1.0	36.4



Chapter from the *Communicable Disease Morbidity Report 1996*, Disease Control Programs.
County of Los Angeles Department of Health Services.

Typhoid fever, carrier	1.2	1.0	1.4	0.6	0.0	0.2	0.6	0.6	0.4	0.2	0.0	0.0	6.2
Typhus fever	0.0	0.4	0.2	0.8	0.8	1.4	1.0	1.2	0.6	1.0	1.0	1.2	9.6
Vibrio ^a	0.0	1.3	0.3	0.8	2.3	3.0	2.0	3.3	2.0	1.3	0.5	0.8	17.3

^aChlamydia, gonorrhea, syphilis and tuberculosis were not included because seasonality may not apply to these diseases.

^bFour-year average (1993-1996)



**Table J. Number of Cases of Selected Notifiable Diseases by Age Group
Los Angeles County, 1996**

Disease	< 1	1-4	5-14	15-34	35-"	45-54	55-64	65 +	Total^a
AIDS	2	3	9	1,011	955	403	117	36	2,536
Amebiasis	1	10	31	76	42	27	12	11	210
Botulism	0	0	0	1	2	1	0	0	4
Brucellosis	0	2	6	1	3	0	0	0	12
Campylobacteriosis	92	310	219	482	224	153	105	146	1,731
Chlamydia	69	10	462	17,468	1,180	272	51	39	20,191
Coccidioidomycosis	0	0	0	17	12	14	15	10	68
Cryptosporidiosis	4	5	9	55	44	22	7	2	148
Cysticercosis	0	1	1	18	5	4	1	0	30
<i>E coli</i> 01 57:H7	0	3	6	2	1	1	1	4	18
Encephalitis	2	3	2	7	7	5	2	5	33
Giardiasis	10	248	202	211	139	68	43	30	951
Gonorrhea	0	25	108	4,446	782	190	44	32	5,733
<i>Haemophilus influenzae</i> type b	0	1	1	2	0	0	0	0	4
Hansen's Disease (Leprosy)	0	0	0	2	2	2	2	1	9
Hepatitis A	9	114	474	392	165	63	51	33	1,301
Hepatitis B	2	0	5	108	58	24	15	18	230
Hepatitis non-A non-B	3	1	3	29	84	57	34	20	231
Hepatitis unspecified	0	2	5	8	5	3	2	3	28
Kawasaki syndrome	0	16	4	0	0	0	0	0	20
Legionellosis	0	0	0	3	0	3	4	2	12
Listeriosis, nonperinatal	0	0	1	3	2	4	2	18	30
Listeriosis, perinatal	0	0	0	5	0	0	0	0	5
Lyme Disease	0	0	0	0	1	0	1	0	2
Malaria	1	3	13	20	7	5	2	8	59
Measles	0	0	1	1	0	0	0	0	2
Meningitis, viral	35	11	29	52	25	15	7	11	185
Meningococcal infections	7	11	7	8	6	7	3	10	59
Mumps	2	7	16	8	3	1	0	0	37
Pertussis	94	15	11	0	0	0	0	0	120
Rheumatic fever, acute	0	0	1	0	0	0	0	1	2
Rubella	0	0	1	1	0	0	0	0	2
Salmonellosis	181	333	267	387	193	141	81	174	1,757
Shigellosis	24	358	262	257	92	53	44	38	1,128
Strongyloidiasis	0	0	0	5	3	0	1	2	11
Syphilis (early latent)	0	1	7	437	225	54	27	5	757
Syphilis (prim. & secon.)	0	0	2	127	57	18	8	4	216
Syphilis (congenital)	126	-	-	-	-	-	-	-	126
'Tetanus	0	0	0	0	0	1	0	0	1
Trichinosis	0	0	0	0	0	0	0	0	0
Tuberculosis	2	49	42	343	270	208	154	307	1,375



Chapter from the *Communicable Disease Morbidity Report 1996*, Disease Control Programs.
County of Los Angeles Department of Health Services.

Tularemia	0	0	0	0	1	0	0	0	1
Typhoid fever, case	0	7	4	13	3	0	1	3	31
Typhoid fever, carrier	0	0	0	0	1	0	0	3	4
Typhus fever	0	0	0	7	4	3	2	0	16
Vibrio	1	0	1	6	7	6	2	0	23

^aTotals include cases with unknown age.

- Not applicable.



**Table K. Incidence Rates of Selected Notifiable Diseases by Age Group
Los Angeles County, 1996**

Disease	Age-group Rates (Cases per 100,000)							
	< 1	1-4	5-14	15-34	35-44	45-54	55-64	65 +
AIDS	1.31	0.41	0.63	38.54	67.56	41.03	18.68	3.90
Amebiasis	0.66	1.36	2.17	2.90	2.97	2.75	1.92	1.19
Botulism	0.00	0.00	0.00	0.04	0.14	0.10	0.00	0.00
Brucellosis	0.00	0.27	0.42	0.04	0.21	0.00	0.00	0.00
Campylobacteriosis	60.43	42.24	15.35	18.37	15.85	15.58	16.77	15.83
Chlamydia	45.30	1.40	32.40	665.60	83.50	27.70	8.20	4.20
Coccidioidomycosis	0.00	0.00	0.00	0.65	0.85	1.43	2.40	1.08
Cryptosporidiosis	2.63	0.68	0.63	2.10	3.11	2.24	1.12	0.22
Cysticercosis	0.00	0.14	0.07	0.69	0.35	0.41	0.16	0.00
<i>E coli</i> 01 57-H7	0.00	0.41	0.42	0.08	0.07	0.10	0.16	0.43
Encephalitis	1.31	0.41	0.14	0.27	0.50	0.51	0.32	0.54
Giardiasis	6.57	33.79	14.16	8.04	9.83	6.92	6.87	3.25
Gonorrhea	0.00	2.80	169.20	55.10	55.10	19.20	7.00	3.50
<i>Haemophilus influenzae</i> type b	0.00	0.14	0.07	0.08	0.00	0.00	0.00	0.00
Hansen's Disease (Leprosy)	0.00	0.00	0.00	0.08	0.14	0.20	0.32	0.11
Hepatitis A	6.57	15.53	33.23	14.94	11.67	6.41	8.14	3.58
Hepatitis B	1.31	0.00	0.35	4.12	4.10	2.44	2.40	1.95
Hepatitis non-A non-B	1.97	0.14	0.21	1.11	5.94	5.80	5.43	2.17
Hepatitis unspecified	0.00	0.27	0.35	0.30	0.35	0.31	0.32	0.33
Kawasaki syndrome	0.00	2.18	0.28	0.00	0.00	0.00	0.00	0.00
Legionellosis	0.00	0.00	0.00	0.11	0.00	0.31	0.64	0.22
Listeriosis, nonperinatal	0.00	0.00	0.07	0.11	0.14	0.41	0.32	1.95
Listeriosis, perinatal'	0.00	0.00	0.00	3.90	0.00	0.00	0.00	0.00
Lyme Disease	0.00	0.00	0.00	0.00	0.07	0.00	0.16	0.00
Malaria	0.66	0.41	0.91	0.76	0.50	0.51	0.32	0.87
Measles	0.00	0.00	0.07	0.04	0.00	0.00	0.00	0.00
Meningitis, viral	22.99	1.50	2.03	1.98	1.77	1.53	1.12	1.19
Meningococcal infections	4.60	1.50	0.49	0.30	0.42	0.71	0.48	1.08
Mumps	1.31	0.95	1.12	0.30	0.21	0.10	0.00	0.00
Pertussis	61.74	2.04	0.77	0.00	0.00	0.00	0.00	0.00
Rheumatic fever, acute	0.00	0.00	0.07	0.00	0.00	0.00	0.00	0.11
Rubella	0.00	0.00	0.07	0.04	0.00	0.00	0.00	0.00
Salmonellosis	118.89	45.38	18.72	14.75	13.65	14.36	12.94	18.86
Shigellosis	15.76	48.78	18.37	9.80	6.51	5.40	7.03	4.12
Strongyloidiasis	0.00	0.00	0.00	0.19	0.21	0.00	0.16	0.22
Syphilis (early latent)	0.00	0.10	0.50	16.70	15.90	5.50	4.30	0.50
Syphilis (prim. & secon.)	0.00	0.00	0.10	4.80	4.00	1.80	1.30	0.40
Syphilis (congenital)	76.80	-	-	-	-	-	-	-
Tetanus	0.00	0.00	0.00	0.00	0.00	0.10	0.00	0.00



Chapter from the *Communicable Disease Morbidity Report 1996*, Disease Control Programs.
County of Los Angeles Department of Health Services.

Trichinosis	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Tuberculosis	1.31	6.68	2.94	13.08	19.10	21.18	24.59	33.28
Tularemia	0.00	0.00	0.00	0.00	0.07	0.00	0.00	0.00
Typhoid fever, case	0.00	0.95	0.28	0.50	0.21	0.00	0.16	0.33
Typhoid fever, carrier	0.00	0.00	0.00	0.00	0.07	0.00	0.00	0.33
Typhus fever	0.00	0.00	0.00	0.27	0.28	0.31	0.32	0.00
Vibrio	0.66	0.00	0.07	0.23	0.50	0.61	0.32	0.00

^aRates for perinatal Listeriosis were **calculated** as cases per 100,000 live births.

-Not applicable.



**Table L. Number of Cases of Selected Notifiable Diseases by Race/Ethnicity
Los Angeles County, 1996**

Disease	Asian	Black	Hispanic	White	Other	Unknown	Total
AIDS	75	567	855	1,039	-	-	2,536
Amebiasis	15	16	106	65	7	1	210
Botulism	0	0	2	2	0	0	4
Brucellosis	0	0	11	1	0	0	12
Campylobacteriosis	132	82	784	717	13	3	1,731
Chlamydia	271	3,170	6,313	969	13	9,455	20,191
Coccidioidomycosis	4	10	25	27	0	2	68
Cryptosporidiosis	2	11	49	57	1	28	148
Cysticercosis	0	1	29	0	0	0	30
<i>E coli</i> 01 57:H7	1	0	1	16	0	0	18
Encephalitis	1	5	10	16	0	1	33
Giardiasis	47	44	458	384	14	4	951
Gonorrhea	38	2,771	997	534	-	1,383	5,723
<i>Haemophilus influenzae</i> type b	0	0	3	1	0	0	4
Hansen's Disease (Leprosy)	3	0	5	1	0	0	9
Hepatitis A	54	55	768	382	9	33	1,301
Hepatitis B	55	46	66	52	1	10	230
Hepatitis non-A non-B	15	46	61	79	2	28	231
Hepatitis unspecified	0	4	21	3	0	0	28
Kawasaki syndrome	12	3	3	1	1	0	20
Legionellosis	0	1	3	7	0	1	12
Listeriosis, nonperinatal	3	5	6	16	0	0	30
Listeriosis, perinatal	0	1	2	2	0	0	5
Lyme Disease	0	0	0	2	0	0	2
Malaria	21	15	17	6	0	0	59
Measles	0	0	0	2	0	0	2
Meningitis, viral	7	28	86	61	0	3	185
Meningococcal infections	0	8	26	23	1	1	59
Mumps	9	1	18	6	2	1	37
Pertussis	4	14	84	18	0	0	120
Rheumatic fever, acute	1	0	0	1	0	0	2
Rubella	0	1	1	0	0	0	2
Salmonellosis	165	202	748	617	25	0	1,757
Shigellosis	25	100	771	222	6	4	1,128
Strongyloidiasis	0	0	11	0	0	0	11
Syphilis (early latent)	6	302	288	68	-	78	742
Syphilis (prim. & secon.)	2	107	60	27	-	20	216
Syphilis (congenital)	1	53	48	7	-	17	126
Tetanus	0	0	1	0	0	0	1
Trichinosis	0	0	0	0	0	0	0
Tuberculosis	377	227	619	148	3	1	1,375
Tularemia	0	0	0	1	0	0	1



Chapter from the *Communicable Disease Morbidity Report 1996*, Disease Control Programs.
County of Los Angeles Department of Health Services.

Typhoid fever, case	9	3	11	5	1	2	31
Typhoid fever, carrier	1	1	1	0	1	0	4
Typhus fever	1	0	3	12	0	0	16
Vibrio	2	0	13	7	0	1	23

^a**Includes** Asian, other, and unknown.



**Table M. Incidence Rate of Selected Notifiable Diseases by Race/Ethnicity
Los Angeles County, 1996**

Disease	Race/Ethnicity Rate (Cases per 100,000)			
	Asian	Black	Hispanic	White
AIDS	6.71'	67.95	21.98	34.19
Amebiasis	1.39	1.92	2.73	2.14
Botulism	0.00	0.00	0.05	0.07
Brucellosis	0.00	0.00	0.28	0.03
Campylobacteriosis	12.22	9.83	20.16	23.60
Chlamydia	47.20	714.90	305.30	60.00
Coccidioidomycosis	0.37	1.20	0.64	0.82
Cryptosporidiosis	0.19	1.32	1.26	1.88
Cysticercosis	0.00	0.12	0.75	0.00
<i>E coli</i> O 157: H7	0.09	0.00	0.03	0.53
Encephalitis	0.09	0.60	0.26	0.53
Giardiasis	4.35	5.27	11.78	12.64
Gonorrhea	4.60	438.10	33.80	23.20
<i>Haemophilus influenzae</i> type b	0.00	0.00	0.08	0.03
Hansen's Disease (Leprosy)	0.28	0.00	0.13	0.03
Hepatitis A	5.00	6.59	19.75	12.57
Hepatitis B	5.09	5.51	1.70	1.71
Hepatitis non-A non-B	1.39	5.51	1.57	2.60
Hepatitis unspecified	0.00	0.48	0.54	0.10
Kawasaki syndrome	1.11	0.36	0.08	0.03
Legionellosis	0.00	0.12	0.08	0.23
Listeriosis, nonperinatal	0.28	0.60	0.15	0.53
Listeriosis, perinatal'	0.00	7.23	2.04	6.76
Lyme Disease	0.00	0.00	0.00	0.07
Malaria	1.94	1.80	0.44	0.20
Measles	0.00	0.00	0.00	0.07
Meningitis, viral	0.65	3.36	2.21	2.01
Meningococcal infections	0.00	0.96	0.67	0.76
Mumps	0.83	0.12	0.46	0.20
Pertussis	0.37	1.68	2.16	0.59
Rheumatic fever, acute	0.09	0.00	0.00	0.03
Rubella	0.00	0.12	0.03	0.00
Salmonellosis	15.27	24.21	19.23	20.30
Shigellosis	2.31	11.98	19.82	7.31
Strongyloidiasis	0.00	0.00	0.28	0.00
Syphilis (early latent)	0.60	40.50	8.30	2.50
Syphilis (prim. & secon.)	0.20	14.10	1.70	1.00
Syphilis (congenital)	11.40	404.30	55.40	22.50
Tetanus	0.00	0.00	0.03	0.00
Trichinosis	0.00	0.00	0.00	0.00
Tuberculosis	34.89	27.20	15.91	4.87



Chapter from the *Communicable Disease Morbidity Report 1996*, Disease Control Programs.
County of Los Angeles Department of Health Services.

Tularemia	0.00	0.00	0.00	0.03
Typhoid fever, case	0.83	0.36	0.28	0.16
Typhoid fever, career	0.09	0.12	0.03	0.00
Typhus fever	0.09	0.00	0.08	0.39
Vibrio	0.19	0.00	0.33	0.23

^aIncludes Asian, other, and unknown.

^bRates for perinatal Listeriosis and congenital syphilis were calculated as cases per 100,000 live births.



**Table N. Number of Cases and Annual Incidence Rate of Selected Notifiable Diseases
by Sex Los Angeles County, 1996**

Disease	Male		Female	
	Cases	Rate (Cases per 100,000)	Cases	Rate (Cases per 100,000)
AIDS	2,282	51.39	254	5.72
Amebiasis	143	3.22	67	1.51
Botulism	2	0.05	2	0.05
Brucellosis	8	0.18	4	0.09
Campylobacteriosis	971	21.87	759	17.10
Chlamydia	4,381	98.70	15,808	356.20
Coccidioidomycosis	53	1.19	15	0.34
Cryptosporidiosis	125	2.81	23	0.52
Cysticercosis	8	0.18	22	0.50
<i>E coli</i> O 157: H7	6	0.14	12	0.27
Encephalitis	23	0.52	10	0.23
Giardiasis	574	12.93	376	8.47
Gonorrhea	3,107	70.00	21616	58.90
<i>Haemophilus influenzae</i> type b	1	0.02	3	0.07
Hansen's Disease (Leprosy)	6	0.14	3	0.07
Hepatitis A	719	16.19	577	13.00
Hepatitis B	147	3.31	83	1.87
Hepatitis non-A non-B	138	3.11	91	2.05
Hepatitis unspecified	14	0.32	14	0.32
Kawasaki syndrome	11	0.25	9	0.20
Legionellosis	10	0.23	2	0.05
Listeriosis, nonperinatal	19	0.43	11	0.25
Listeriosis, perinatal'	0	0.00	5	5.76
Lyme Disease	2	0.05	0	0.00
Malaria	32	0.72	27	0.61
Measles	1	0.02	1	0.02
Meningitis, viral	102	2.30	83	1.87
Meningococcal infections	30	0.68	29	0.65
Mumps	21	0.47	16	0.36
Pertussis	55	1.24	65	1.46
Rheumatic fever, acute	1	0.02	1	0.02
Rubella	2	0.05	0	0.00
Salmonellosis	858	19.32	897	20.21
Shigellosis	549	12.36	579	13.04
Strongyloidiasis	7	0.16	4	0.09
Syphilis (early latent)	375	8.50	366	8.30
Syphilis (prim. & secon.)	129	2.90	86	1.90
Syphilis (congenital)	N/A	N/A	N/A	N/A
Tetanus	1	0.02	0	0.00
Trichinosis	0	0.00	0	0.00



Chapter from the *Communicable Disease Morbidity Report 1996*, Disease Control Programs.
County of Los Angeles Department of Health Services.

Tuberculosis	905	20.38	470	10.59
Tularemia	1	0.02	0	0.00
Typhoid fever, case	13	0.29	18	0.41
Typhoid fever, career	2	0.05	2	0.05
Typhus fever	7	0.16	9	0.20
Vibrio	19	0.43	4	0.09

^aRates for perinatal Listeriosis were calculated as cases per 100.000 live births



Table 0-1. Selected Notifiable Diseases
Antelope Valley, San Fernando, and West Valley Health Districts
Los Angeles County, 1996

Disease	Frequency			Rate (Cases per 100,000)		
	Antelope	San Fern	W Valley	Antelope	San Fern	W Valley
AIDS	34	18	114	12.47	5.07	15.42
Amebiasis	2	3	8	0.73	0.85	1.08
Botulism	0	0	0	0.00	0.00	0.00
Brucellosis	0	0	0	0.00	0.00	0.00
Campylobacteriosis	39	107	140	14.30	30.15	18.94
Chlamydia ^a	N/A	750	917	N/A	139.30	144.60
Coccidioidomycosis	7	7	10	2.57	1.97	1.35
Cryptosporidiosis	2	5	4	0.73	1.41	0.54
Cysticercosis	1	1	5	0.37	0.28	0.68
<i>E coli</i> O 157: H7	0	3	0	0.00	0.85	0.00
Encephalitis	0	1	1	0.00	0.28	0.14
Foodborne outbreaks	0	0	1	0.00	0.00	0.14
Giardiasis	27	66	56	9.90	18.60	7.57
Gonorrhea ^c	N/A	176	210	N/A	31.90	32.40
<i>Haemophilus influenzae</i> type b	0	0	1	0.00	0.00	0.14
Hansen's Disease (leprosy)	0	0	0	0.00	0.00	0.00
Hepatitis A	27	27	60	9.90	7.61	8.12
Hepatitis B	14	4	12	5.14	1.13	1.62
Hepatitis non-A non-B	27	10	8	9.90	2.82	1.08
Hepatitis unspecified	1	0	2	0.37	0.00	0.27
Kawasaki syndrome	0	0	4	0.00	0.00	0.54
Legionellosis	0	0	2	0.00	0.00	0.27
Listeriosis, nonperinatal	1	1	0	0.37	0.28	0.00
Listeriosis, perinatal ^b	0	0	0	0.00	0.00	0.00
Lyme Disease	0	0	0	0.00	0.00	0.00
Malaria	0	2	4	0.00	0.56	0.54
Measles	0	0	0	0.00	0.00	0.00
Meningitis, viral	7	3	15	2.57	0.85	2.03
Meningococcal infections	1	1	4	0.37	0.28	0.54
Mumps	0	1	2	0.00	0.28	0.27
Pertussis	4	3	10	1.47	0.85	1.35
Rheumatic fever, acute	0	0	0	0.00	0.00	0.00
Rubella	0	0	0	0.00	0.00	0.00
Salmonellosis	48	65	136	17.61	18.32	18.40
Shigellosis	13	23	57	4.77	6.48	7.71
Strongyloidiasis	0	0	1	0.00	0.00	0.14
Syphilis (early latent) ^c	N/A	14	25	N/A	2.30	3.40
Syphilis (prim. & secon.) ^c	N/A	6	7	N/A	1.00	1.00
Syphilis (congenital) ^c	N/A	7	7	N/A	N/A	N/A
Tetanus	0	0	0	0.00	0.00	0.00



Chapter from the *Communicable Disease Morbidity Report 1996*, Disease Control Programs.
County of Los Angeles Department of Health Services.

Trichinosis	0	0	0	0.00	0.00	0.00
Tuberculosis	24	20	75	8.80	5.64	10.15
Tularemia	0	0	0	0.00	0.00	0.00
Typhoid fever, case	0	0	3	0.00	0.00	0.41
Typhoid fever, carrier	0	0	1	0.00	0.00	0.14
Typhus fever	0	0	0	0.00	0.00	0.00
Vibrio	0	0	1	0.00	0.00	0.14

^aData for Antelope Valley Health District were included in San Fernando Health District.

^bRates for perinatal Listeriosis were calculated as cases per 100.000 women aged 15 to 44 years.



**Table 0-2. Selected Notifiable Diseases
East Valley and Glendale Health Districts
Los Angeles County, 1996**

Disease	Frequency		Rate (Cases per 100,000)	
	E Valley	Glendale	E Valley	Glendale
AIDS	121	31	29.55	9.60
Amebiasis	10	3	2.44	0.93
Botulism	0	0	0.00	0.00
Brucellosis	1	1	0.24	0.31
Campylobacteriosis	94	53	22.96	16.41
Chlamydia	858	319	244.20	115.10
Coccidioidomycosis	2	0	0.49	0.00
Cryptosporidiosis	9	2	2.20	0.62
Cysticercosis	3	1	0.73	0.31
<i>E. coli</i> 01 57:H7	3	0	0.73	0.00
Encephalitis	3	1	0.73	0.31
Foodborne outbreaks	0	1	0.00	0.31
Giardiasis	62	14	15.14	4.33
Gonorrhea	144	54	40.10	19.0
<i>Haemophilus influenzae</i> type b	2	0	0.49	0.00
Hansen's Disease (Leprosy)	1	0	0.24	0.00
Hepatitis A	50	15	12.21	4.64
Hepatitis B	4	2	0.98	0.62
Hepatitis non-A non-B	9	6	2.20	1.86
Hepatitis unspecified	3	0	0.73	0.00
Kawasaki syndrome	0	1	0.00	0.31
Legionellosis	0	0	0.00	0.00
Listeriosis, nonperinatal	0	1	0.00	0.31
Listeriosis, perinatal ^a	0	0	0.00	0.00
Lyme Disease	0	0	0.00	0.00
Malaria	6	0	1.47	0.00
Measles	2	0	0.49	0.00
Meningitis, viral	10	9	2.44	2.79
Meningococcal infections	5	2	1.22	0.62
Mumps	2	0	0.49	0.00
Pertussis	0	3	0.00	0.93
Rheumatic fever, acute	0	0	0.00	0.00
Rubella	0	0	0.00	0.00
Salmonellosis	85	59	20.76	18.26
Shigellosis	53	20	12.94	6.19
Strongyloidiasis	0	0	0.00	0.00
Syphilis (early latent)	12	7	3.00	2.20
Syphilis (prim. & secon.)	6	3	1.50	0.90
Syphilis (congenital)	4	2	N/A	N/A
Tetanus	0	0	0.00	0.00
Trichinosis	0	0	0.00	0.00



Chapter from the *Communicable Disease Morbidity Report 1996*, Disease Control Programs.
County of Los Angeles Department of Health Services.

Tuberculosis	55	40	13.43	12.38
Tularemia	0	0	0.00	0.00
Typhoid fever, case	2	0	0.49	0.00
Typhoid fever, carrier	0	0	0.00	0.00
Typhus fever	0	2	0.00	0.62
Vibrio	2	0	0.49	0.00

^aRates for perinatal Listeriosis were calculated as cases per 100.000 women aged 15 to 44 years.



**Table 0-4. Selected Notifiable Diseases
Pomona Health District
Los Angeles County, 1996**

Disease	Frequency	Rate (Cases per 100,000)
	Pomona	Pomona
AIDS	32	6.04
Amebiasis	4	0.76
Botulism	0	0.00
Brucellosis	0	0.00
Campylobacteriosis	62	11.71
Chlamydia	760	167.30
Coccidioidomycosis	4	0.76
Cryptosporidiosis	1	0.19
Cysticercosis	1	0.19
E coli O157:H7	0	0.00
Encephalitis	2	0.38
Foodborne outbreaks	0	0.00
Giardiasis	20	3.78
Gonorrhea	134	28.80
<i>Haemophilus influenzae</i> type b	0	0.00
Hansen's Disease (Leprosy)	1	0.19
Hepatitis A	76	14.35
Hepatitis B	11	2.08
Hepatitis non-A non-B	3	0.57
Hepatitis unspecified	0	0.00
Kawasaki syndrome	2	0.38
Legionellosis	0	0.00
Listeriosis, nonperinatal	0	0.00
Listeriosis, perinatal ^a	0	0.00
Lyme Disease	0	0.00
Malaria	3	0.57
Measles	0	0.00
Meningitis, viral	15	2.83
Meningococcal infections	2	0.38
Mumps	2	0.38
Pertussis	7	1.32
Rheumatic fever, acute	0	0.00
Rubella	0	0.00
Salmonellosis	88	16.62
Shigellosis	45	8.50
Strongyloidiasis	0	0.00
Syphilis (early latent)	11	2.10
Syphilis (prim. & secon.)	1	0.20
Syphilis (congenital)	1	N/A
Tetanus	0	0.00



Chapter from the *Communicable Disease Morbidity Report 1996*, Disease Control Programs.
County of Los Angeles Department of Health Services.

Trichinosis	0	0.00
Tuberculosis	61	11.52
Tularemia	0	0.00
Typhoid fever, case	2	0.38
Typhoid fever, career	0	0.00
Typhus fever	0	0.00
Vibrio		0.19

^aRates for perinatal Listeriosis were calculated as cases per 100.000 women aged 15 to 44 years.



**Table 0-3. Selected Notifiable Diseases
Alhambra, El Monte and Foothill Health Districts
Los Angeles County, 1996**

Disease	Frequency			Rate (Cases per 100,000)		
	Alhambra	El Monte	Foothill	Alhambra	El Monte	Foothill
AIDS	19	34	25	5.10	7.22	8.35
Amebiasis	7	16	4	1.88	3.40	1.34
Botulism	0	0	0	0.00	0.00	0.00
Brucellosis	0	1	0	0.00	0.21	0.00
Campylobacteriosis	47	70	37	12.62	14.86	12.36
Chlamydia	373	822	359	116.80	203.40	139.70
Coccidioidomycosis	2	0	1	0.54	0.00	0.33
Cryptosporidiosis	3	3	2	0.81	0.64	0.67
Cysticercosis	2	0	1	0.54	0.00	0.33
<i>E coli</i> 01 57:H7	1	0	0	0.27	0.00	0.00
Encephalitis	0	1	1	0.00	0.21	0.33
Foodborne outbreaks	0	0	0	0.00	0.00	0.00
Giardiasis	37	61	34	9.94	12.95	11.35
Gonorrhea	41	79	72	12.50	19.10	27.4
<i>Haemophilus influenzae</i> type b	1	0	0	0.27	0.00	0.00
Hansen's Disease (Leprosy)	0	0	0	0.00	0.00	0.00
Hepatitis A	31	80	17	8.33	16.99	5.68
Hepatitis B	26	11	1	6.98	2.34	0.33
Hepatitis non-A non-B	11	8	2	2.95	1.70	0.67
Hepatitis unspecified	0	0	1	0.00	0.00	0.33
Kawasaki syndrome	1	1	1	0.27	0.21	0.33
Legionellosis	1	0	2	0.27	0.00	0.67
Listeriosis, nonperinatal	3	3	1	0.81	0.64	0.33
Listeriosis, perinatal'	0	0	1	0.00	0.00	1.52
Lyme Disease	0	0	0	0.00	0.00	0.00
Malaria	3	1	2	0.81	0.21	0.67
Measles	0	0	0	0.00	0.00	0.00
Meningitis, viral	4	14	10	1.07	2.97	3.34
Meningococcal infections	1	1	4	0.27	0.21	1.34
Mumps	3	3	1	0.81	0.64	0.33
Pertussis	0	6	2	0.00	1.27	0.67
Rheumatic fever, acute	0	0	0	0.00	0.00	0.00
Rubella	0	0	0	0.00	0.00	0.00
Salmonellosis	55	68	65	14.77	14.44	21.71
Shigellosis	27	65	21	7.25	13.80	7.01
Strongyloidiasis	0	0	0	0.00	0.00	0.00
Syphilis (early latent)	8	14	10	2.20	3.00	3.40
Syphilis (prim. & secon.)	1	4	3	0.30	0.90	1.00
Syphilis (congenital)	0	3	1	N/A	N/A	N/A
Tetanus	0	0	0	0.00	0.00	0.00



Chapter from the *Communicable Disease Morbidity Report 1996*, Disease Control Programs.
County of Los Angeles Department of Health Services.

Trichinosis	0	0	0	0.00	0.00	0.00
Tuberculosis	71	82	26	19.07	17.41	8.68
Tularemia	0	0	0	0.00	0.00	0.00
Typhoid fever, case	2	3	0	0.54	0.64	0.00
Typhoid fever, career	1	0	0	0.27	0.00	0.00
Typhus fever	4	0	4	1.07	0.00	1.34
Vibrio	0	4	0	0.00	0.85	0.00

^aRates for perinatal Listeriosis were calculated as cases per 100.000 women aged 15 to 44 years.



**Table 0-5. Selected Notifiable Diseases
Bellflower, East LA and Whittier Health Districts
Los Angeles County, 1996**

Disease	Frequency			Rate (Cases per 100,000)		
	Bellflower	East LA	Whittier	Bellflower	East LA	Whittier
AIDS	41	33	38	11.84	14.39	12.00
Amebiasis	4	2	2	1.16	0.87	0.63
Botulism	0	0	0	0.00	0.00	0.00
Brucellosis	0	1	1	0.00	0.44	0.32
Campylobacteriosis	60	48	59	17.33	20.93	18.63
Chlamydia	550	400	470	185.20	203.30	173.00
Coccidioidomycosis	6	0	1	1.73	0.00	0.32
Cryptosporidiosis	2	2	2	0.58	0.87	0.63
Cysticercosis	1	0	0	0.29	0.00	0.00
<i>E coli</i> 01 57:H7	0	0	1	0.00	0.00	0.32
Encephalitis	4	1	1	1.16	0.44	0.32
Foodborne outbreaks	0	0	0	0.00	0.00	0.00
Giardiasis	23	18	18	6.64	7.85	5.68
Gonorrhea	95	43	76	31.30	21.40	27.30
<i>Haemophilus influenzae</i> type b	0	0	0	0.00	0.00	0.00
Hansen's Disease (Leprosy)	0	1	0	0.00	0.44	0.00
Hepatitis A	67	41	41	19.35	17.88	12.95
Hepatitis B	6	3	1	1.73	1.31	0.32
Hepatitis non-A non-B	2	6	0	0.58	2.62	0.00
Hepatitis unspecified	1	1	0	0.29	0.44	0.00
Kawasaki syndrome	0	1	0	0.00	0.44	0.00
Legionellosis	0	0	1	0.00	0.00	0.32
Listeriosis, nonperinatal	0	0	1	0.00	0.00	0.32
Listeriosis, perinatal ^a	0	0	0	0.00	0.00	0.00
Lyme Disease	0	0	0	0.00	0.00	0.00
Malaria	4	0	0	1.16	0.00	0.00
Measles	0	0	0	0.00	0.00	0.00
Meningitis, viral	7	3	3	2.02	1.31	0.95
Meningococcal infections	6	1	1	1.73	0.44	0.32
Mumps	4	0	0	1.16	0.00	0.00
Pertussis	4	2	2	1.16	0.87	0.63
Rheumatic fever, acute	0	0	0	0.00	0.00	0.00
Rubella	0	0	0	0.00	0.00	0.00
Salmonellosis	74	30	66	21.38	13.08	20.84
Shigellosis	30	41	44	8.67	17.88	13.89
Strongyloidiasis	1	0	0	0.29	0.00	0.00
Syphilis (early latent)	12	15	13	3.50	6.70	4.20
Syphilis (prim. & secon.)	4	5	1	1.20	2.20	0.30
Syphilis (congenital)	1	4	1	N/A	N/A	N/A
Tetanus	0	0	0	0.00	0.00	0.00



Chapter from the *Communicable Disease Morbidity Report 1996*, Disease Control Programs.
County of Los Angeles Department of Health Services.

Trichinosis	0	0	0	0.00	0.00	0.00
Tuberculosis	42	33	17	12.13	14.39	5.37
Tularemia	0	0	0	0.00	0.00	0.00
Typhoid fever, case	1	2	0	0.29	0.87	0.00
Typhoid fever, carrier	0	0	0	0.00	0.00	0.00
Typhus fever	0	0	0	0.00	0.00	0.00
Vibrio	1	1	0	0.29	0.44	0.00

^aRates for perinatal Listeriosis were calculated as cases per 100.000 women aged 15 to 44 years.



**Table 0-6. Selected Notifiable Diseases
Central and Northeast Health Districts
Los Angeles County, 1996**

Disease	Frequency		Rate (Cases per 100,000)	
	Central	NE	Central	NE
AIDS	209	83	63.98	23.08
Amebiasis	14	6	4.29	1.67
Botulism	0	0	0.00	0.00
Brucellosis	0	3	0.00	0.83
Campylobacteriosis	46	88	14.08	24.47
Chlamydia	799	690	285.20	223.60
Coccidioidomycosis	4	0	1.22	0.00
Cryptosporidiosis	17	8	5.20	2.22
Cysticercosis	0	1	0.00	0.28
<i>E coli</i> 01 57:H7	0	0	0.00	0.00
Encephalitis	0	2	0.00	0.56
Foodborne outbreaks	0	0	0.00	0.00
Giardiasis	25	33	7.66	9.17
Gonorrhea	302	95	105.30	30.10
<i>Haemophilus influenzae</i> type b	0	0	0.00	0.00
Hansen's Disease (Leprosy)	1	0	0.31	0.00
Hepatitis A	55	63	16.84	17.52
Hepatitis B	12	11	3.67	3.06
Hepatitis non-A non-B	1	5	0.31	1.39
Hepatitis unspecified	0	4	0.00	1.11
Kawasaki syndrome	1	0	0.31	0.00
Legionellosis	0	0	0.00	0.00
Listeriosis, nonperinatal	0	1	0.00	0.28
Listeriosis, perinatal ^a	0	1	0.00	1.21
Lyme Disease	0	0	0.00	0.00
Malaria	2	1	0.61	0.28
Measles	0	0	0.00	0.00
Meningitis, viral	4	9	1.22	2.50
Meningococcal infections	0	3	0.00	0.83
Mumps	0	2	0.00	0.56
Pertussis	11	7	3.37	1.95
Rheumatic fever, acute	0	1	0.00	0.28
Rubella	0	0	0.00	0.00
Salmonellosis	39	76	11.94	21.13
Shigellosis	69	78	21.12	21.69
Strongyloidiasis	1	0	0.31	0.00
Syphilis (early latent)	106	22	33.00	6.20
Syphilis (prim. & secon.)	30	2	9.30	0.60
Syphilis (congenital)	13	3	N/A	N/A
Tetanus	0	0	0.00	0.00



Chapter from the *Communicable Disease Morbidity Report 1996*, Disease Control Programs.
County of Los Angeles Department of Health Services.

Trichinosis	0	0	0.00	0.00
Tuberculosis	190	75	58.17	20.85
Tularemia	1	0	0.31	0.00
Typhoid fever, case	1	2	0.31	0.56
Typhoid fever, carrier	0	0	0.00	0.00
Typhus fever	0	6	0.00	1.67
Vibrio	0	1	0.00	0.28

^aRates for perinatal Listeriosis were calculated as cases per 100.000 women aged 15 to 44 years.



**Table 0-7. Selected Notifiable Diseases
Hollywood-Wilshire and West Health Districts
Los Angeles County, 1996**

Disease	Frequency		Rate (Cases per 100,000)	
	Hol-Wil	West	Hol-Wil	West
AIDS	507	114	102.26	18.89
Amebiasis	49	15	9.88	2.49
Botulism	0	0	0.00	0.00
Brucellosis	0	0	0.00	0.00
Campylobacteriosis	90	132	18.15	21.87
Chlamydia	1,011	610	237.60	117.80
Coccidioidomycosis	1	2	0.20	0.33
Cryptosporidiosis	39	14	7.87	2.32
Cysticercosis	3	0	0.61	0.00
<i>E coli</i> 01 57:H7	1	5	0.20	0.83
Encephalitis	2	3	0.40	0.50
Foodborne outbreaks	0	4	0.00	0.66
Giardiasis	57	58	11.50	9.61
Gonorrhea	498	186	114.40	35.10
<i>Haemophilus influenzae</i> type b	0	0	0.00	0.00
Hansen's Disease (Leprosy)	2	0	0.40	0.00
Hepatitis A	110	42	22.19	6.96
Hepatitis B	28	10	5.65	1.66
Hepatitis non-A non-B	52	1	10.49	0.17
Hepatitis unspecified	1	1	0.20	0.17
Kawasaki syndrome	1	0	0.20	0.00
Legionellosis	2	0	0.40	0.00
Listeriosis, nonperinatal	2	8	0.40	1.33
Listeriosis, perinatal'	0	1	0.00	0.71
Lyme Disease	0	0	0.00	0.00
Malaria	8	5	1.61	0.83
Measles	0	0	0.00	0.00
Meningitis, viral	13	4	2.62	0.66
Meningococcal infections	1	2	0.20	0.33
Mumps	3	0	0.61	0.00
Pertussis	5	3	1.01	0.50
Rheumatic fever, acute	0	0	0.00	0.00
Rubella	1	1	0.20	0.17
Salmonellosis	106	102	21.38	16.90
Shigellosis	126	38	25.41	6.30
Strongyloidiasis	3	0	0.61	0.00
Syphilis (early latent)	75	15	15.40	2.50
Syphilis (prim. & secon.)	28	1	5.70	0.20
Syphilis (congenital)	4	1	N/A	N/A
Tetanus	0	0	0.00	0.00



Chapter from the *Communicable Disease Morbidity Report 1996*, Disease Control Programs.
County of Los Angeles Department of Health Services.

Trichinosis	0	0	0.00	0.00
Tuberculosis	111	52	22.39	8.62
Tularemia	0	0	0.00	0.00
Typhoid fever, case	4	3	0.81	0.50
Typhoid fever, carrier	1	1	0.20	0.17
Typhus fever	0	0	0.00	0.00
Vibrio	1	1	0.20	0.17

^aRates for perinatal Listeriosis were calculated as cases per 100.000 women aged 15 to 44 years.



**Table 0-8. Selected Notifiable Diseases
Compton, San Antonio and South Health Districts
Los Angeles County, 1996**

Disease	Frequency			Rate (Cases per 100,000)		
	Comp	S Antonio	South	Comp	S Antonio	South
AIDS	37	50	44	13.38	11.62	26.47
Amebiasis	5	13	2	1.81	3.02	1.20
Botulism	0	1	0	0.00	0.23	0.00
Brucellosis	1	0	1	0.36	0.00	0.60
Campylobacteriosis	33	69	28	11.93	16.03	16.84
Chlamydia	1,077	815	926	453.90	220.70	650.80
Coccidioidomycosis	2	1	3	0.72	0.23	1.80
Cryptosporidiosis	3	2	2	1.08	0.46	1.20
Cysticercosis	1	4	0	0.36	0.93	0.00
<i>E. coli</i> O 157: H7	0	0	0	0.00	0.00	0.00
Encephalitis	0	0	0	0.00	0.00	0.00
Foodborne outbreaks	0	0	0	0.00	0.00	0.00
Giardiasis	27	59	18	9.76	13.71	10.83
Gonorrhea	457	110	451	188.20	29.10	309.10
<i>Haemophilus influenzae</i> type b	0	0	0	0.00	0.00	0.00
Hansen's Disease (Leprosy)	0	2	0	0.00	0.46	0.00
Hepatitis A	53	75	34	19.16	17.42	20.45
Hepatitis B	10	9	8	3.62	2.09	4.81
Hepatitis non-A non-B	7	7	33	2.53	1.63	19.85
Hepatitis unspecified	2	4	1	0.72	0.93	0.60
Kawasaki syndrome	1	1	0	0.36	0.23	0.00
Legionellosis	0	1	0	0.00	0.23	0.00
Listeriosis, nonperinatal	0	2	0	0.00	0.46	0.00
Listeriosis, perinatal'	0	0	0	0.00	0.00	0.00
Lyme Disease	0	0	0	0.00	0.00	0.00
Malaria	1	0	0	0.36	0.00	0.00
Measles	0	0	0	0.00	0.00	0.00
Meningitis, viral	8	8	4	2.89	1.86	2.41
Meningococcal injections	5	2	0	1.81	0.46	0.00
Mumps	1	1	1	0.36	0.23	0.60
Pertussis	3	10	4	1.08	2.32	2.41
Rheumatic fever, acute	0	0	0	0.00	0.00	0.00
Rubella	0	0	0	0.00	0.00	0.00
Salmonellosis	45	99	32	16.27	23.00	19.25
Shigellosis	41	73	43	14.83	16.96	25.87
Strongyloidiasis	0	0	0	0.00	0.00	0.00
Syphilis (early latent)	53	33	72	19.50	7.80	44.10
Syphilis (prim. & secon.)	17	7	32	6.20	1.60	19.40
Syphilis (congenital)	4	3	18	N/A	N/A	N/A
Tetanus	0	0	1	0.00	0.00	0.60



Chapter from the *Communicable Disease Morbidity Report 1996*, Disease Control Programs.
County of Los Angeles Department of Health Services.

Trichinosis	0	0	0	0.00	0.00	0.00
Tuberculosis	49	51	32	17.72	11.85	19.25
Tularemia	0	0	0	0.00	0.00	0.00
Typhoid fever, case	0	0	0	0.00	0.00	0.00
Typhoid fever, carrier	0	0	0	0.00	0.00	0.00
Typhus fever	0	0	0	0.00	0.00	0.00
Vibrio	1	2		0.36	0.46	0.00

^aRates for perinatal Listeriosis were calculated as cases per 100.000 women aged 15 to 44 years.



**Table 0-9. Selected Notifiable Diseases
Southeast, Southwest, and Inglewood Health Districts
Los Angeles County, 1996**

Disease	Frequency			Rate (Cases per 100,000)		
	Ing	SE	SW	Ing	SE	SW
AIDS	69	36	99	17.27	22.92	26.68
Amebiasis	5	3	5	1.25	1.91	1.35
Botulism	2	0	0	0.50	0.00	0.00
Brucellosis	1	0	0	0.25	0.00	0.00
Campylobacteriosis	54	27	68	13.52	17.19	18.32
Chlamydia	1,653	679	1,626	482.20	504.00	510.70
Coccidioidomycosis	3	2	2	0.75	1.27	0.54
Cryptosporidiosis	1	1	5	0.25	0.64	1.35
Cysticercosis	1	0	1	0.25	0.00	0.27
E coli 01 57:H7	0	0	0	0.00	0.00	0.00
Encephalitis	2	2	1	0.50	1.27	0.27
Foodborne outbreaks	1	0	0	0.25	0.00	0.00
Giardiasis	27	17	16	6.76	10.82	4.31
Gonorrhea	770	184	651	219.50	133.50	199.80
<i>Haemophilus influenzae</i> type b	0	0	0	0.00	0.00	0.00
Hansen's Disease (Leprosy)	0	0	0	0.00	0.00	0.00
Hepatitis A	45	28	24	11.26	17.83	6.47
Hepatitis B	4	6	12	1.00	3.82	3.23
Hepatitis non-A non-B	4	3	8	1.00	1.91	2.16
Hepatitis unspecified	1	0	0	0.25	0.00	0.00
Kawasaki syndrome	1	0	1	0.25	0.00	0.27
Legionellosis	0	0	1	0.00	0.00	0.27
Listeriosis, nonperinatal	2	0	2	0.50	0.00	0.54
Listeriosis, perinatal ^a	0	1	1	0.00	2.47	1.14
Lyme Disease	0	0	0	0.00	0.00	0.00
Malaria	4	0	5	1.00	0.00	1.35
Measles	0	0	0	0.00	0.00	0.00
Meningitis, viral	5	2	7	1.25	1.27	1.89
Meningococcal infections	3	5	3	0.75	3.18	0.81
Mumps	2	1	1	0.50	0.64	0.27
Pertussis	8	6	12	2.00	3.82	3.23
Rheumatic fever, acute	0	0	0	0.00	0.00	0.00
Rubella	0	0	0	0.00	0.00	0.00
Salmonellosis	100	29	65	25.03	18.47	17.51
Shigellosis	38	31	46	9.51	19.74	12.39
Strongyloidiasis	0	3	1	0.00	1.91	0.27
Syphilis (early latent)	81	37	80	20.60	24.00	21.90
Syphilis (prim. & secon.)	18	11	19	4.50	7.10	5.20
Syphilis (congenital)	12	8	26	N/A	N/A	N/A
Tetanus	0	0	0	0.00	0.00	0.00



Chapter from the *Communicable Disease Morbidity Report 1996*, Disease Control Programs.
County of Los Angeles Department of Health Services.

Trichinosis	0	0	0	0.00	0.00	0.00
Tuberculosis	69	44	96	17.27	28.02	25.87
Tularemia	0	0	0	0.00	0.00	0.00
Typhoid fever, case	1	0	1	0.25	0.00	0.27
Typhoid fever, career	0	0	0	0.00	0.00	0.00
Typhus fever	0	0	0	0.00	0.00	0.00
Vibrio	0	1	0	0.00	0.64	0.00

^aRates for perinatal Listeriosis were calculated as cases per 100,000 women aged 15 to 44 years.

N/A - Not available



**Table 0-10. Selected Notifiable Diseases
Harbor and Torrance Health Districts
Los Angeles County, 1996**

Disease	Frequency		Rate (Cases per 100,000)	
	Harbor	Torrance	Harbor	Torrance
AIDS	22	44	11.17	10.17
Amebiasis	4	7	2.03	1.62
Botulism	0	0	0.00	0.00
Brucellosis	0	0	0.00	0.00
Campylobacteriosis	83	113	42.14	26.12
Chlamydia	386	474	228.40	127.70
Coccidioidomycosis	0	1	0.00	0.23
Cryptosporidiosis	2	4	1.02	0.92
Cysticercosis	0	0	0.00	0.00
<i>E coli</i> 01 57:H7	1	3	0.51	0.69
Encephalitis	0	1	0.00	0.23
Foodborne outbreaks	0	1	0.00	0.23
Giardiasis	34	84	17.26	19.42
Gonorrhea	72	125	41.60	32.90
<i>Haemophilus influenzae</i> type b	0	0	0.00	0.00
Hansen's Disease, (Leprosy)	0	1	0.00	0.23
Hepatitis A	53	83	26.91	19.19
Hepatitis B	1	7	0.51	1.62
Hepatitis non-A non-B	0	1	0.00	0.23
Hepatitis unspecified	1	1	0.51	0.23
Kawasaki syndrome	0	1	0.00	0.23
Legionellosis	0	2	0.00	0.46
Listeriosis, nonperinatal	0	2	0.00	0.46
Listeriosis, perinatal ^a	0	0	0.00	0.00
Lyme Disease	0	2	0.00	0.46
Malaria	0	3	0.00	0.69
Measles	0	0	0.00	0.00
Meningitis, viral	5	8	2.54	1.85
Meningococcal infections	1	3	0.51	0.69
Mumps	1	3	0.51	0.69
Pertussis	4	4	2.03	0.92
Rheumatic fever, acute	0	1	0.00	0.23
Rubella	0	0	0.00	0.00
Salmonellosis	34	98	17.26	22.65
Shigellosis	19	43	9.65	9.94
Strongyloidiasis	0	0	0.00	0.00
Syphilis (early latent)	9	7	4.70	1.60
Syphilis (prim. & secon.)	2	6	1.00	1.40
Syphilis (congenital)	1	1	N/A	N/A
Tetanus	0	0	0.00	0.00



Chapter from the *Communicable Disease Morbidity Report 1996*, Disease Control Programs.
County of Los Angeles Department of Health Services.

Trichinosis	0	0	0.00	0.00
Tuberculosis	12	42	6.09	9.71
Tularemia	0	0	0.00	0.00
Typhoid fever, case	1	0	0.51	0.00
Typhoid fever, carrier	0	0	0.00	0.00
Typhus fever	0	0	0.00	0.00
Vibrio	1	2	0.51	0.46

^aRates for perinatal Listeriosis were calculated as cases per 100,000 women aged 15 to 44 years.

N/A - Not available