

**Table G. Reported Cases of Selected Notifiable Diseases by Year of Onset
Los Angeles County, 1992-1997**

Disease	Year of Onset						Previous 5-year Average	5-Yr 95% Upper Limit ^a
	1992	1993	1994	1995	1996	1997		
AIDS ^b	3,689	3,515	3,306	2,985	2,249	1,560	3,149	4,261
Amebiasis	239	290	230	187	215	148	232	306
Botulism	0	1	3	2	4	3	2	5
Brucellosis	6	4	11	3	12	6	7	15
Campylobacteriosis	1,396	1,385	1,301	1,401	1,736	1,523	1,444	1,774
Chlamydia ^c	21,679	20,521	21,258	18,659	20,191	23,021	20,462	22,748
Cholera	3	3	4	2	0	0	2	5
Coccidioidomycosis	98	106	98	76	69	40	89	121
Cryptosporidiosis	N/A	102	231	211	149	77	N/A	N/A
Cysticercosis	43	59	41	26	33	34	40	65
Dengue	0	0	0	2	0	2	0	2
E. coli O157:H7	N/A	9	13	9	18	20	N/A	N/A
Encephalitis	35	59	36	59	33	34	44	71
Foodborne outbreaks ^c	1	7	12	15	12	42	9	20
Giardiasis	1,622	1,616	1,105	940	971	770	1,251	1,921
Gonorrhea	14,001	11,080	8,802	7,807	5,723	5,825	9,483	15,710
<i>Haemophilus influenzae</i> type b	38	22	20	6	4	10	18	45
Hansen's Disease (Leprosy)	17	24	25	10	9	18	17	32
Hepatitis A ^c	1,196	1,040	1,193	1,062	1,371	1,480	1,172	1,432
Hepatitis B	1,077	288	289	231	247	109	426	1,141
Hepatitis C	255	148	115	205	246	23	194	313
Hepatitis unspecified	45	32	40	22	28	16	33	51
Kawasaki syndrome	47	49	46	39	20	20	40	64
Legionellosis ^c	16	11	11	16	12	32	13	18
Listeriosis, nonperinatal	23	25	26	26	30	14	26	31
Listeriosis, perinatal	22	23	11	10	5	8	14	30
Lyme disease	5	0	2	5	3	4	3	7
Malaria	69	40	40	76	62	55	57	90
Measles	37	51	16	7	2	4	23	63
Meningitis, viral	936	490	250	167	185	227	406	1,039
Meningococcal infections	85	98	91	52	59	74	77	117
Mumps	57	48	51	42	37	39	47	62
Pertussis	140	104	58	103	120	32	105	164
Psittacosis	1	1	0	0	0	1	0	1
Q-fever	0	0	0	0	0	0	0	0
Relapsing fever	1	1	0	0	0	0	0	1
Rheumatic fever, acute	3	3	5	1	2	1	3	6
Rubella	6	3	4	3	3	5	4	6
Salmonellosis	1,617	1,565	2,091	2,084	1,773	1,675	1,826	2,317
Shigellosis	1,882	1,532	1,336	1,747	1,130	848	1,525	2,120
Strongyloidiasis	45	42	24	17	11	4	28	57
Syphilis (early latent)	2,050	1,660	1,024	926	744	648	1,281	2,361
Syphilis (prim. & secon.)	726	511	326	271	216	105	410	819
Syphilis (congenital)	322	257	202	161	126	76	214	366
Tetanus	1	1	0	4	1	4	1	4
Trichinosis	2	0	2	1	0	2	1	3
Tuberculosis	2,198	1,940	1,794	1,622	1,375	1,347	1,786	2,397
Tularemia	1	0	0	1	1	2	1	2
Typhoid fever, case	45	42	38	26	31	26	36	52
Typhoid fever, carrier	8	8	5	6	4	1	6	10
Typhus fever	0	14	8	11	16	13	10	22
Vibrio	N/A	14	13	12	23	28	N/A	N/A

^a The normal distribution assumption may not apply to some rare diseases.

^b All data are adjusted for report delay and the 1993 AIDS case definition change.

^c 1997 data over 95% upper limit.

**Table H. Annual Incidence Rates of Selected Notifiable Diseases by Year of Onset
Los Angeles County, 1992-1997**

Disease	Annual Incidence Rate (Cases per 100,000)					
	1992	1993	1994	1995	1996	1997
AIDS ^a	42.88	39.96	38.19	34.10	25.33	17.24
Amebiasis	2.78	3.30	2.66	2.14	2.42	1.64
Botulism	0.00	0.01	0.03	0.02	0.05	0.03
Brucellosis	0.07	0.05	0.13	0.03	0.14	0.07
Campylobacteriosis	16.23	15.75	15.03	16.00	19.55	16.83
Chlamydia	252.01	233.31	245.57	213.15	227.40	254.30
Cholera	0.03	0.03	0.05	0.02	0.00	0.00
Coccidioidomycosis	1.14	1.21	1.13	0.87	0.78	0.44
Cryptosporidiosis	N/A	1.16	2.67	2.41	1.68	0.85
Cysticercosis	0.50	0.67	0.47	0.30	0.37	0.38
Dengue	0.00	0.00	0.00	0.02	0.00	0.02
E. coli O157:H7	N/A	0.10	0.15	0.10	0.20	0.22
Encephalitis	0.41	0.67	0.42	0.67	0.37	0.38
Giardiasis	18.86	18.37	12.76	10.74	10.93	8.51
Gonorrhea	162.76	125.97	101.68	89.18	64.50	64.40
<i>Haemophilus influenzae</i> type b	0.44	0.25	0.23	0.07	0.05	0.11
Hansen's disease (Leprosy)	0.20	0.27	0.29	0.11	0.10	0.20
Hepatitis A	13.90	11.82	13.78	12.13	14.99	16.40
Hepatitis B	12.52	3.27	3.34	2.64	2.78	1.20
Hepatitis C	2.96	1.68	1.33	2.34	2.77	0.25
Hepatitis unspecified	0.52	0.36	0.46	0.25	0.32	0.18
Kawasaki syndrome	0.55	0.56	0.53	0.45	0.23	0.22
Legionellosis	0.19	0.13	0.13	0.18	0.14	0.35
Listeriosis, nonperinatal	0.27	0.28	0.30	0.30	0.34	0.15
Listeriosis, perinatal ^b	11.98	13.03	6.01	5.81	3.19	5.02
Lyme disease	0.06	0.00	0.02	0.06	0.03	0.04
Malaria	0.80	0.45	0.46	0.87	0.70	0.61
Measles	0.43	0.58	0.18	0.08	0.02	0.04
Meningitis, viral	10.88	5.57	2.89	1.91	2.08	2.51
Meningococcal infections	0.99	1.11	1.05	0.59	0.66	0.82
Mumps	0.66	0.55	0.59	0.48	0.42	0.43
Pertussis	1.63	1.18	0.67	1.18	1.35	0.35
Psittacosis	0.01	0.01	0.00	0.00	0.00	0.01
Q-fever	0.00	0.00	0.00	0.00	0.00	0.00
Relapsing fever	0.01	0.01	0.00	0.00	0.00	0.00
Rheumatic fever, acute	0.03	0.03	0.06	0.01	0.02	0.01
Rubella	0.07	0.03	0.05	0.03	0.03	0.06
Salmonellosis	18.80	17.79	24.16	23.81	19.97	18.51
Shigellosis	21.88	17.42	15.43	19.96	12.73	9.37
Strongyloidiasis	0.52	0.48	0.28	0.19	0.12	0.04
Syphilis (early latent)	23.83	18.87	11.83	10.58	8.40	7.20
Syphilis (prim. & secon.)	8.44	5.81	3.77	3.10	2.40	1.20
Syphilis (congenital) ^b	175.47	145.70	125.60	98.10	76.80	48.10
Tetanus	0.01	0.01	0.00	0.05	0.01	0.04
Trichinosis	0.02	0.00	0.02	0.01	0.00	0.02
Tuberculosis	25.55	22.06	20.72	18.53	15.48	14.88
Tularemia	0.01	0.00	0.00	0.01	0.01	0.02
Typhoid fever, case	0.52	0.48	0.44	0.30	0.35	0.29
Typhoid fever, carrier	0.09	0.09	0.06	0.07	0.05	0.01
Typhus fever	0.00	0.16	0.09	0.13	0.18	0.14
Vibrio	N/A	0.16	0.15	0.14	0.26	0.31

^aAll data are adjusted for report delay and the 1993 AIDS case definition change.

^bRates for perinatal listeriosis and congenital syphilis were calculated as cases per 100,000 live births.

**Table I. Five-Year Average of Notifiable Diseases by Month of Onset
Los Angeles County, 1993-1997**

Disease	Jan	Feb	Mar	Apr	May	June	July	Aug	Sept	Oct	Nov	Dec	Total ^a
AIDS ^b	253.0	221.0	272.0	237.0	232.0	230.0	224.0	213.0	220.0	217.0	190.0	203.0	2723.0
Amebiasis	23.8	17.2	21.8	19.0	17.6	13.6	19.8	20.2	18.0	15.8	13.2	14.0	214.0
Botulism	0.2	0.2	0.0	0.0	0.4	0.0	0.4	0.4	0.2	0.2	0.0	0.4	2.4
Brucellosis	1.2	0.4	0.2	0.8	0.2	1.0	1.2	0.6	0.4	0.6	0.2	0.4	7.2
Campylobacteriosis	106.2	90.2	92.8	119.8	157.6	161.2	160.2	141.6	135.6	119.4	104.4	80.2	1469.2
Cholera	0.4	0.2	0.2	0.4	0.0	0.2	0.2	0.0	0.0	0.0	0.0	0.0	1.6
Coccidioidomycosis	15.0	9.0	5.0	7.0	5.0	4.6	6.0	4.6	4.6	5.0	5.4	6.0	77.2
Cryptosporidiosis	35.0	9.6	11.8	6.2	8.6	9.0	10.6	10.0	19.8	13.4	9.6	10.4	154.0
Cysticercosis	7.6	3.0	3.4	2.6	3.0	3.6	2.8	2.6	3.0	2.2	3.2	1.6	38.6
Dengue	0.2	0.0	0.0	0.2	0.0	0.4	0.0	0.0	0.0	0.0	0.0	0.0	0.8
E. coli O157:H7	0.6	0.8	1.0	1.0	1.0	1.8	0.6	1.2	1.4	2.2	1.4	0.8	13.8
Encephalitis	11.4	2.8	4.8	3.4	4.4	3.6	2.0	2.2	2.0	1.8	3.4	2.6	44.4
Giardiasis	115.0	69.2	95.0	76.6	88.6	83.6	97.6	117.8	116.0	95.8	65.8	59.2	1080.2
<i>Haemophilus influenzae</i> type b	2.0	3.0	1.0	0.6	0.2	1.0	0.4	0.8	0.6	1.0	1.2	0.6	12.4
Hansen's disease (Leprosy)	2.6	1.2	1.6	0.8	2.0	2.6	0.6	1.4	0.2	0.2	1.0	0.8	15.0
Hepatitis A	219.8	83.4	84.6	74.0	84.2	81.0	83.6	103.8	118.6	114.8	102.4	84.2	1234.4
Hepatitis B	24.8	18.2	17.8	19.2	16.8	18.8	19.2	16.4	20.2	19.4	15.2	16.4	222.4
Hepatitis C	12.6	10.6	11.8	14.0	12.2	10.8	11.4	10.0	9.4	9.2	8.6	11.8	132.4
Hepatitis unspecified	8.6	1.8	2.2	1.8	1.6	2.4	2.4	1.2	2.0	1.4	1.0	1.2	27.6
Kawasaki syndrome	4.2	4.0	4.0	4.2	2.8	1.4	2.6	1.8	2.0	2.4	1.8	2.8	34.0
Legionellosis	0.6	0.8	2.0	1.0	0.6	0.4	1.4	1.6	1.2	1.6	3.2	1.2	15.6
Listeriosis, nonperinatal	1.2	2.0	1.6	0.8	1.4	2.4	2.4	4.2	2.0	2.8	1.4	1.6	23.8
Listeriosis, perinatal	1.2	0.4	1.0	0.8	1.2	1.0	1.2	1.0	1.4	1.0	0.8	0.2	11.2
Lyme disease	0.6	0.0	0.2	0.0	0.4	0.0	0.6	0.2	0.4	0.2	0.0	0.0	2.6
Malaria	5.0	3.4	2.8	4.0	7.0	4.2	4.8	7.2	6.6	3.4	3.4	3.0	54.8
Measles	0.6	0.2	0.4	1.4	4.2	4.6	3.2	0.6	0.2	0.2	0.0	0.2	15.8
Meningitis, viral	18.6	11.2	16.2	17.8	22.6	33.0	33.8	34.8	27.4	23.0	15.8	9.6	263.8
Meningococcal infections	13.0	11.8	6.8	4.8	5.6	5.0	3.4	4.0	2.0	2.4	2.4	7.8	69.0
Mumps	6.4	4.6	4.8	4.2	6.0	3.2	2.4	2.6	4.2	2.6	2.2	0.2	43.4
Pertussis	5.6	4.6	5.8	7.2	7.6	7.8	8.4	10.0	8.0	8.0	5.8	4.6	83.4
Psittacosis	0.0	0.0	0.0	0.2	0.0	0.2	0.0	0.0	0.0	0.0	0.0	0.0	0.4
Relapsing fever	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.2	0.0	0.0	0.2
Rheumatic fever, acute	0.6	0.2	0.0	0.4	0.2	0.0	0.0	0.2	0.4	0.4	0.0	0.0	2.4
Rubella	0.0	0.0	0.8	0.2	0.2	0.8	0.2	0.2	0.2	0.8	0.2	0.0	3.6
Salmonellosis	120.4	98.6	133.6	129.0	161.6	161.6	189.2	229.0	203.4	181.6	135.6	94.4	1838.0
Shigellosis	89.6	62.4	66.8	66.2	78.8	117.8	158.4	207.8	196.2	133.2	85.8	55.4	1318.4
Strongyloidiasis	1.6	1.4	1.2	2.8	2.2	2.2	1.0	2.2	1.8	1.0	1.2	1.0	19.6
Tetanus	0.4	0.2	0.0	0.4	0.4	0.0	0.2	0.0	0.2	0.0	0.0	0.2	2.0
Tularemia	0.0	0.0	0.0	0.0	0.2	0.0	0.4	0.0	0.0	0.2	0.0	0.0	0.8
Typhoid fever, case	3.2	2.6	2.8	2.0	4.2	3.0	3.8	4.0	3.4	1.4	0.6	1.0	32.0
Typhoid fever, carrier	1.4	0.6	0.0	0.2	0.0	0.8	0.6	0.6	0.2	0.6	0.2	0.2	5.4
Typhus fever	0.4	0.4	0.2	0.8	0.8	1.6	1.4	1.4	0.8	2.0	1.4	1.2	12.4
Vibrio	0.2	1.4	0.2	0.6	1.8	2.4	2.2	2.8	2.6	2.0	0.4	0.6	17.2

^a Chlamydia, gonorrhea, syphilis and tuberculosis were not included because seasonality may not apply to these diseases.

^b Month of diagnosis is unknown for 54 cases.

**Table J. Number of Cases of Selected Notifiable Diseases by Age Group
Los Angeles County, 1997**

Disease	<1	1-4	5-14	15-34	35-44	45-54	55-64	65+	Total ^a
AIDS	3	2	4	624	597	226	84	20	1,560
Amebiasis	0	11	17	55	28	22	10	5	148
Botulism	0	0	0	0	2	1	0	0	3
Brucellosis	0	0	2	1	1	2	0	0	6
Campylobacteriosis	81	270	213	413	188	143	86	128	1,523
Chlamydia	51	8	434	20,206	1,382	280	58	36	23,021
Coccidioidomycosis	0	0	2	9	12	7	4	6	40
Cryptosporidiosis	1	7	3	23	23	15	3	1	77
Cysticercosis	0	2	6	16	7	0	2	1	34
Dengue	0	0	0	0	2	0	0	0	2
E. coli O157:H7	0	5	8	2	1	0	1	3	20
Encephalitis	1	3	7	6	5	3	1	8	34
Giardiasis	5	169	190	183	102	66	33	22	770
Gonorrhea	13	4	88	4,440	860	197	66	23	5,825
<i>Haemophilus influenzae</i> type	3	3	0	0	0	0	0	4	10
Hansen's disease (Leprosy)	0	0	0	3	5	4	5	1	18
Hepatitis A	2	113	520	477	184	111	31	41	1,480
Hepatitis B	0	0	4	64	20	9	2	9	109
Hepatitis C	0	0	0	5	12	4	1	1	23
Hepatitis unspecified	0	1	5	4	2	1	0	3	16
Kawasaki syndrome	6	11	3	0	0	0	0	0	20
Legionellosis	0	0	1	2	3	6	8	12	32
Listeriosis, nonperinatal	0	0	1	2	0	0	2	9	14
Listeriosis, perinatal	0	0	0	6	1	1	0	0	8
Lyme disease	0	0	0	0	1	1	2	0	4
Malaria	1	3	7	20	16	2	4	2	55
Measles	1	1	1	1	0	0	0	0	4
Meningitis, viral	59	21	55	55	22	3	3	9	227
Meningococcal infections	17	10	7	17	4	5	3	11	74
Mumps	1	8	17	8	1	2	1	1	39
Pertussis	27	5	0	0	0	0	0	0	32
Psittacosis	0	0	0	0	0	0	1	0	1
Relapsing fever	0	0	0	0	0	0	0	0	0
Rheumatic fever, acute	0	1	0	0	0	0	0	0	1
Rubella	0	0	0	0	5	0	0	0	5
Salmonellosis	149	291	263	414	182	136	82	147	1,675
Shigellosis	17	254	197	193	82	41	29	33	848
Strongyloidiasis	0	0	0	1	2	0	0	1	4
Syphilis (early latent)	0	0	2	375	182	59	20	8	648
Syphilis (prim. & secon.)	0	0	0	52	41	9	3	0	105
Syphilis (congenital)	76	-	-	-	-	-	-	-	76
Tetanus	0	0	0	0	1	1	1	1	4
Trichinosis	0	0	2	0	0	0	0	0	2
Tuberculosis	3	47	37	361	253	192	158	296	1,347
Tularemia	0	0	0	0	1	1	0	0	2
Typhoid fever, case	0	3	8	10	3	1	0	1	26
Typhoid fever, carrier	0	0	0	0	0	0	0	1	1
Typhus fever	0	0	2	1	4	5	1	0	13
Vibrio	1	0	1	15	6	0	2	3	28

^aTotals include cases with unknown age.

- Not applicable.

**Table K. Incidence Rates of Selected Notifiable Diseases by Age Group
Los Angeles County, 1997**

Disease	Age-group Rates (Cases per 100,000)							
	<1	1-4	5-14	15-34	35-44	45-54	55-64	65+
AIDS	1.90	0.31	0.27	23.73	40.70	22.19	13.13	1.37
Amebiasis	0.00	1.73	1.16	2.09	1.84	2.07	1.52	0.54
Botulism	0.00	0.00	0.00	0.00	0.13	0.09	0.00	0.00
Brucellosis	0.00	0.00	0.14	0.04	0.07	0.19	0.00	0.00
Campylobacteriosis	51.24	42.44	14.58	15.70	12.35	13.48	13.11	13.79
Chlamydia	22.10	2.50	29.70	767.90	90.80	26.40	8.80	3.90
Coccidioidomycosis	0.00	0.00	0.14	0.34	0.79	0.66	0.61	0.65
Cryptosporidiosis	0.63	1.10	0.21	0.87	1.51	1.41	0.46	0.11
Cysticercosis	0.00	0.31	0.41	0.61	0.46	0.00	0.30	0.11
Dengue	0.00	0.00	0.00	0.00	0.13	0.00	0.00	0.00
E. coli O157:H7	0.00	0.79	0.55	0.08	0.07	0.00	0.15	0.32
Encephalitis	0.63	0.47	0.48	0.23	0.33	0.28	0.15	0.86
Giardiasis	3.16	26.56	13.01	6.96	6.70	6.22	5.03	2.37
Gonorrhea	8.20	0.60	6.00	168.80	56.50	18.60	10.10	2.50
<i>Haemophilus influenzae</i> type b	1.90	0.47	0.00	0.00	0.00	0.00	0.00	0.43
Hansen's disease (Leprosy)	0.00	0.00	0.00	0.11	0.33	0.38	0.76	0.11
Hepatitis A	1.27	17.76	35.60	18.14	12.09	10.46	4.73	4.42
Hepatitis B	0.00	0.00	0.27	2.43	1.31	0.85	0.30	0.97
Hepatitis C	0.00	0.00	0.00	0.19	0.79	0.38	0.15	0.11
Hepatitis unspecified	0.00	0.16	0.34	0.15	0.13	0.09	0.00	0.32
Kawasaki syndrome	3.80	1.73	0.21	0.00	0.00	0.00	0.00	0.00
Legionellosis	0.00	0.00	0.07	0.08	0.20	0.57	1.22	1.29
Listeriosis, nonperinatal	0.00	0.00	0.07	0.08	0.00	0.00	0.30	0.97
Listeriosis, perinatal ^a	0.00	0.00	0.00	4.63	3.47	189.75	0.00	0.00
Lyme disease	0.00	0.00	0.00	0.00	0.07	0.09	0.30	0.00
Malaria	0.63	0.47	0.48	0.76	1.05	0.19	0.61	0.22
Measles	0.63	0.16	0.07	0.04	0.00	0.00	0.00	0.00
Meningitis, viral	37.32	3.30	3.77	2.09	1.45	0.28	0.46	0.97
Meningococcal infections	10.75	1.57	0.48	0.65	0.26	0.47	0.46	1.19
Mumps	0.63	1.26	1.16	0.30	0.07	0.19	0.15	0.11
Pertussis	17.08	0.79	0.00	0.00	0.00	0.00	0.00	0.00
Psittacosis	0.00	0.00	0.00	0.00	0.00	0.00	0.15	0.00
Relapsing fever	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Rheumatic fever, acute	0.00	0.16	0.00	0.00	0.00	0.00	0.00	0.00
Rubella	0.00	0.00	0.00	0.00	0.33	0.00	0.00	0.00
Salmonellosis	94.26	45.74	18.00	15.74	11.96	12.82	12.50	15.84
Shigellosis	10.75	39.92	13.49	7.34	5.39	3.86	4.42	3.56
Strongyloidiasis	0.00	0.00	0.00	0.04	0.13	0.00	0.00	0.11
Syphilis (early latent)	0.00	0.00	0.13	14.30	12.00	5.60	3.00	0.86
Syphilis (prim. & secon.)	0.00	0.00	0.00	2.00	2.70	0.80	0.50	0.00
Syphilis (congenital)	48.10	-	-	-	-	-	-	-
Tetanus	0.00	0.00	0.00	0.00	0.07	0.09	0.15	0.11
Trichinosis	0.00	0.00	0.14	0.00	0.00	0.00	0.00	0.00
Tuberculosis	1.90	7.39	2.53	13.73	16.63	18.10	24.09	31.89
Tularemia	0.00	0.00	0.00	0.00	0.07	0.09	0.00	0.00
Typhoid fever, case	0.00	0.47	0.55	0.38	0.20	0.09	0.00	0.11
Typhoid fever, carrier	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.11
Typhus fever	0.00	0.00	0.14	0.04	0.26	0.47	0.15	0.00
Vibrio	0.63	0.00	0.07	0.57	0.39	0.00	0.30	0.32

^aRates for perinatal listeriosis were calculated as cases per 100,000 live births.

-Not applicable.

**Table L. Number of Cases of Selected Notifiable Diseases by Race/Ethnicity
Los Angeles County, 1997**

Disease	Asian	Black	Hispanic	White	Other	Unknown	
AIDS	41	399	651	463	4	2	1,560
Amebiasis	14	11	83	38	2	0	148
Botulism	0	0	0	3	0	0	3
Brucellosis	0	0	5	1	0	0	6
Campylobacteriosis	125	54	688	632	8	16	1,523
Chlamydia	378	3,717	7,459	1,248	32	10,187	23,021
Coccidioidomycosis	7	10	8	12	0	3	40
Cryptosporidiosis	0	6	23	29	0	19	77
Cysticercosis	1	0	29	3	0	1	34
Dengue	0	0	2	0	0	0	2
E. coli O157:H7	3	2	0	15	0	0	20
Encephalitis	2	4	13	13	0	2	34
Giardiasis	43	38	388	289	6	6	770
Gonorrhea	48	2,529	863	509	0	1,876	5,825
<i>Haemophilus influenzae</i> type b	0	3	3	4	0	0	10
Hansen's disease (Leprosy)	9	0	8	0	0	1	18
Hepatitis A	43	91	855	436	11	44	1,480
Hepatitis B	20	18	33	36	0	2	109
Hepatitis C	0	3	14	6	0	0	23
Hepatitis unspecified	2	1	9	4	0	0	16
Kawasaki syndrome	6	2	6	4	0	2	20
Legionellosis	0	5	6	21	0	0	32
Listeriosis, nonperinatal	0	2	4	8	0	0	14
Listeriosis, perinatal	0	1	2	5	0	0	8
Lyme disease	0	0	0	4	0	0	4
Malaria	14	12	16	10	0	3	55
Measles	1	1	2	0	0	0	4
Meningitis, viral	13	20	117	68	4	5	227
Meningococcal infections	1	9	38	21	0	5	74
Mumps	4	1	23	11	0	0	39
Pertussis	0	3	23	6	0	0	32
Psittacosis	0	0	0	0	0	1	1
Relapsing fever	0	0	0	0	0	0	0
Rheumatic fever, acute	0	0	1	0	0	0	1
Rubella	1	0	0	4	0	0	5
Salmonellosis	203	199	655	582	8	28	1,675
Shigellosis	28	44	586	170	8	12	848
Strongyloidiasis	0	0	3	0	0	1	4
Syphilis (early latent)	4	258	243	38	0	105	648
Syphilis (prim. & secon.)	0	56	32	8	0	9	105
Syphilis (congenital)	0	28	46	2	0	0	76
Tetanus	0	0	4	0	0	0	4
Trichinosis	0	1	1	0	0	0	2
Tuberculosis	429	187	597	132	2	0	1,347
Tularemia	0	0	1	0	0	1	2
Typhoid fever, case	7	1	13	2	0	3	26
Typhoid fever, carrier	0	0	0	0	0	1	1
Typhus fever	1	0	2	10	0	0	13
Vibrio	5	1	8	14	0	0	28

**Table M. Incidence Rate of Selected Notifiable Diseases by Race/Ethnicity
Los Angeles County, 1997**

Disease	Race/Ethnicity Rate (Cases per 100,000)			
	Asian	Black	Hispanic	White
AIDS	3.56	48.40	16.12	15.49
Amebiasis	1.21	1.33	2.05	1.27
Botulism	0.00	0.00	0.00	0.10
Brucellosis	0.00	0.00	0.12	0.03
Campylobacteriosis	10.84	6.55	17.03	21.14
Chlamydia	58.80	808.60	331.10	74.90
Coccidioidomycosis	0.61	1.21	0.20	0.40
Cryptosporidiosis	0.00	0.73	0.57	0.97
Cysticercosis	0.09	0.00	0.72	0.10
Dengue	0.00	0.00	0.05	0.00
E. coli O157:H7	0.26	0.24	0.00	0.50
Encephalitis	0.17	0.49	0.32	0.43
Giardiasis	3.73	4.61	9.60	9.67
Gonorrhea	6.10	452.50	31.50	25.10
<i>Haemophilus influenzae</i> type b	0.00	0.36	0.07	0.13
Hansen's Disease (Leprosy)	0.78	0.00	0.20	0.00
Hepatitis A	3.73	11.04	21.17	14.58
Hepatitis B	1.73	2.18	0.82	1.20
Hepatitis C	0.00	0.36	0.35	0.20
Hepatitis unspecified	0.17	0.12	0.22	0.13
Kawasaki syndrome	0.52	0.24	0.15	0.13
Legionellosis	0.00	0.61	0.15	0.70
Listeriosis, nonperinatal	0.00	0.24	0.10	0.27
Listeriosis, perinatal ^a	0.00	0.12	0.05	0.17
Lyme Disease	0.00	0.00	0.00	0.13
Malaria	1.21	1.46	0.40	0.33
Measles	0.09	0.12	0.05	0.00
Meningitis, viral	1.13	2.43	2.90	2.27
Meningococcal infections	0.09	1.09	0.94	0.70
Mumps	0.35	0.12	0.57	0.37
Pertussis	0.00	0.36	0.57	0.20
Psittacosis	0.00	0.00	0.00	0.00
Relapsing fever	0.00	0.00	0.00	0.00
Rheumatic fever, acute	0.00	0.00	0.02	0.00
Rubella	0.09	0.00	0.00	0.13
Salmonellosis	17.61	24.14	16.21	19.47
Shigellosis	2.43	5.34	14.51	5.69
Strongyloidiasis	0.00	0.00	0.07	0.00
Syphilis (early latent)	0.40	37.40	7.20	1.50
Syphilis (prim. & secon.)	0.00	7.40	0.90	0.30
Syphilis (congenital) ^a	0.00	241.80	44.20	6.90
Tetanus	0.00	0.00	0.10	0.00
Trichinosis	0.00	0.12	0.02	0.00
Tuberculosis	37.21	22.68	14.78	4.42
Tularemia	0.00	0.00	0.02	0.00
Typhoid fever, case	0.61	0.12	0.32	0.07
Typhoid fever, carrier	0.00	0.00	0.00	0.00
Typhus fever	0.09	0.00	0.05	0.33
Vibrio	0.43	0.12	0.20	0.47

^a Rates for perinatal listeriosis and congenital syphilis were calculated as cases per 100,000 live births.

**Table N. Number of Cases and Annual Incidence Rate of Selected Notifiable Diseases by Sex
Los Angeles County, 1997**

Disease	Male		Female	
	Cases	Rate (Cases per 100,000)	Cases	Rate (Cases per 100,000)
AIDS	1,362	30.21	198	4.36
Amebiasis	88	1.95	60	1.32
Botulism	1	0.02	2	0.04
Brucellosis	4	0.09	2	0.04
Campylobacteriosis	834	18.50	689	15.17
Chlamydia	5,327	118.10	17,694	389.50
Coccidioidomycosis	30	0.67	10	0.22
Cryptosporidiosis	59	1.31	18	0.40
Cysticercosis	13	0.29	21	0.46
Dengue	0	0.00	2	0.04
E. coli O157:H7	8	0.18	12	0.26
Encephalitis	17	0.38	17	0.37
Giardiasis	419	9.29	350	7.70
Gonorrhea	3,172	70.40	2,650	58.30
<i>Haemophilus influenzae</i> type b	6	0.13	4	0.09
Hansen's disease (Leprosy)	11	0.29	2	0.11
Hepatitis A	907	20.12	573	12.61
Hepatitis B	70	1.55	39	0.86
Hepatitis C	15	0.33	8	0.18
Hepatitis unspecified	5	0.11	11	0.24
Kawasaki syndrome	8	0.18	12	0.26
Legionellosis	20	0.44	12	0.26
Listeriosis, nonperinatal	8	0.18	6	0.13
Listeriosis, perinatal ^a	3	0.07	3	0.07
Lyme disease	3	0.07	1	0.02
Malaria	31	0.69	23	0.51
Measles	3	0.07	1	0.02
Meningitis, viral	119	2.64	108	2.38
Meningococcal infections	35	0.78	39	0.86
Mumps	18	0.40	21	0.46
Pertussis	9	0.19	23	0.51
Psittacosis	0	0.00	1	0.02
Relapsing fever	0	0.00	0	0.00
Rheumatic fever, acute	0	0.00	1	0.02
Rubella	2	0.04	3	0.07
Salmonellosis	832	18.45	841	18.51
Shigellosis	409	9.07	439	9.66
Strongyloidiasis	2	0.04	2	0.04
Syphilis (early latent)	328	7.30	317	7.00
Syphilis (prim. & secon.)	73	1.60	32	0.70
Syphilis (congenital)	N/A	N/A	N/A	N/A
Tetanus	3	0.07	1	0.02
Trichinosis	1	0.02	1	0.02
Tuberculosis	858	19.03	489	10.76
Tularemia	1	0.02	1	0.02
Typhoid fever, case	19	0.42	7	0.15
Typhoid fever, carrier	0	0.00	1	0.02
Typhus fever	9	0.20	4	0.09
Vibrio	19	0.42	9	0.20

^aRates for perinatal listeriosis were calculated as cases per 100,000 live births.

**Table O-1. Selected Notifiable Diseases
Antelope Valley, San Fernando, and West Valley Health Districts
Los Angeles County, 1997**

Disease	Frequency			Rate (Cases per 100,000)		
	Antelope	San Fern	W Valley	Antelope	San Fern	W Valley
AIDS	25	21	94	8.34	5.59	12.55
Amebiasis	2	3	7	0.67	0.80	0.93
Botulism	0	1	0	0.00	0.27	0.00
Brucellosis	0	0	0	0.00	0.00	0.00
Campylobacteriosis	32	97	107	10.68	25.83	14.28
Chlamydia ^a	N/A	1,087	1,470	N/A	164.50	196.20
Coccidioidomycosis	6	2	10	2.00	0.53	1.33
Cryptosporidiosis	0	1	5	0.00	0.27	0.67
Cysticercosis	0	3	3	0.00	0.80	0.40
Dengue	0	0	0	0.00	0.00	0.00
E. coli O157:H7	0	4	4	0.00	1.07	0.53
Encephalitis	2	0	4	0.67	0.00	0.53
Giardiasis	23	29	56	7.67	7.72	7.48
Gonorrhea ^a	N/A	204	180	N/A	35.90	28.60
<i>Haemophilus influenzae</i> type b	0	0	1	0.00	0.00	0.13
Hansen's disease (Leprosy)	0	2	1	0.00	0.53	0.13
Hepatitis A	33	44	111	11.01	11.72	14.82
Hepatitis B	4	2	3	1.33	0.53	0.40
Hepatitis C	3	1	0	1.00	0.27	0.00
Hepatitis unspecified	2	0	3	0.67	0.00	0.40
Kawasaki syndrome	0	3	0	0.00	0.80	0.00
Legionellosis	0	1	2	0.00	0.27	0.27
Listeriosis, nonperinatal	0	0	0	0.00	0.00	0.00
Listeriosis, perinatal ^b	0	0	1	0.00	0.00	0.60
Lyme disease	0	0	0	0.00	0.00	0.00
Malaria	1	1	5	0.33	0.27	0.67
Measles	0	0	0	0.00	0.00	0.00
Meningitis, viral	6	7	21	2.00	1.86	2.80
Meningococcal infections	3	1	5	1.00	0.27	0.67
Mumps	0	0	1	0.00	0.00	0.13
Pertussis	0	1	1	0.00	0.27	0.13
Psittacosis	0	0	0	0.00	0.00	0.00
Relapsing fever	0	0	0	0.00	0.00	0.00
Rheumatic fever, acute	0	0	0	0.00	0.00	0.00
Rubella	0	0	1	0.00	0.00	0.13
Salmonellosis	36	73	143	12.01	19.44	19.09
Shigellosis	5	21	61	1.67	5.59	8.14
Strongyloidiasis	0	0	0	0.00	0.00	0.00
Syphilis (early latent) ^a	N/A	17	13	N/A	2.50	1.70
Syphilis (prim. & secon.) ^a	N/A	3	0	N/A	0.40	0.00
Syphilis (congenital) ^a	N/A	4	4	N/A	N/A	N/A
Tetanus	0	2	0	0.00	0.53	0.00
Trichinosis	0	0	0	0.00	0.00	0.00
Tuberculosis	20	34	88	6.67	9.05	11.75
Tularemia	0	1	0	0.00	0.27	0.00
Typhoid fever, case	0	0	2	0.00	0.00	0.27
Typhoid fever, carrier	0	0	0	0.00	0.00	0.00
Typhus fever	0	1	0	0.00	0.27	0.00
Vibrio	0	0	3	0.00	0.00	0.40

^aData for Antelope Valley Health District were included in San Fernando Health District.

^bRates for perinatal listeriosis were calculated as cases per 100,000 women aged 15 to 44 years.

**Table O-2. Selected Notifiable Diseases
East Valley and Glendale Health Districts
Los Angeles County, 1997**

Disease	Frequency		Rate (Cases per 100,000)	
	E Valley	Glendale	E Valley	Glendale
AIDS	84	31	19.84	8.97
Amebiasis	12	6	2.84	1.74
Botulism	0	0	0.00	0.00
Brucellosis	2	0	0.47	0.00
Campylobacteriosis	84	57	19.85	16.50
Chlamydia	1,059	429	250.30	124.20
Coccidioidomycosis	2	1	0.47	0.29
Cryptosporidiosis	9	3	2.13	0.87
Cysticercosis	2	0	0.47	0.00
Dengue	0	1	0.00	0.29
E. coli O157:H7	1	3	0.24	0.87
Encephalitis	3	2	0.71	0.58
Giardiasis	42	12	9.93	3.47
Gonorrhea	117	48	32.80	16.50
<i>Haemophilus influenzae</i> type b	0	0	0.00	0.00
Hansen's disease (Leprosy)	0	0	0.00	0.00
Hepatitis A	106	26	25.05	7.53
Hepatitis B	1	1	0.24	0.29
Hepatitis C	2	1	0.47	0.29
Hepatitis unspecified	1	0	0.24	0.00
Kawasaki syndrome	0	0	0.00	0.00
Legionellosis	0	1	0.00	0.29
Listeriosis, nonperinatal	0	0	0.00	0.00
Listeriosis, perinatal ^a	0	0	0.00	0.00
Lyme disease	0	0	0.00	0.00
Malaria	2	2	0.47	0.58
Measles	0	0	0.00	0.00
Meningitis, viral	6	7	1.42	2.03
Meningococcal infections	1	3	0.24	0.87
Mumps	2	3	0.47	0.87
Pertussis	3	2	0.71	0.58
Psittacosis	0	0	0.00	0.00
Relapsing fever	0	0	0.00	0.00
Rheumatic fever, acute	0	0	0.00	0.00
Rubella	0	0	0.00	0.00
Salmonellosis	58	57	13.71	16.50
Shigellosis	42	13	9.93	3.76
Strongyloidiasis	0	0	0.00	0.00
Syphilis (early latent)	11	11	2.60	3.20
Syphilis (prim. & secon.)	4	2	0.90	0.60
Syphilis (congenital)	1	0	N/A	N/A
Tetanus	0	0	0.00	0.00
Trichinosis	0	0	0.00	0.00
Tuberculosis	25	72	5.91	20.84
Tularemia	0	0	0.00	0.00
Typhoid fever, case	0	0	0.00	0.00
Typhoid fever, carrier	0	0	0.00	0.00
Typhus fever	0	1	0.00	0.29
Vibrio	1	2	0.24	0.58

^aRates for perinatal listeriosis were calculated as cases per 100,000 women aged 15 to 44 years.

**Table O-3. Selected Notifiable Diseases
Alhambra, El Monte and Foothill Health Districts
Los Angeles County, 1997**

Disease	Frequency			Rate (Cases per 100,000)		
	Alhambra	El Monte	Foothill	Alhambra	El Monte	Foothill
AIDS	84	25	25	5.11	5.24	7.94
Amebiasis	6	11	6	1.53	2.30	1.91
Botulism	0	0	0	0.00	0.00	0.00
Brucellosis	0	0	1	0.00	0.00	0.32
Campylobacteriosis	56	59	25	14.30	12.36	7.94
Chlamydia	369	817	358	114.90	208.70	138.50
Coccidioidomycosis	2	0	1	0.51	0.00	0.32
Cryptosporidiosis	0	2	1	0.00	0.42	0.32
Cysticercosis	0	2	2	0.00	0.42	0.64
Dengue	0	0	0	0.00	0.00	0.00
E. coli O157:H7	2	0	0	0.51	0.00	0.00
Encephalitis	0	2	2	0.00	0.42	0.64
Giardiasis	18	42	21	4.60	8.80	6.67
Gonorrhea	32	79	66	9.70	19.70	24.80
<i>Haemophilus influenzae</i> type b	0	1	0	0.00	0.21	0.00
Hansen's disease (Leprosy)	1	2	0	0.26	0.42	0.00
Hepatitis A	29	78	28	7.40	16.34	8.90
Hepatitis B	8	6	5	2.04	1.26	1.59
Hepatitis C	0	2	0	0.00	0.42	0.00
Hepatitis unspecified	0	3	0	0.00	0.63	0.00
Kawasaki syndrome	0	0	1	0.00	0.00	0.32
Legionellosis	0	0	0	0.00	0.00	0.00
Listeriosis, nonperinatal	2	2	0	0.51	0.42	0.00
Listeriosis, perinatal ^a	0	0	0	0.00	0.00	0.00
Lyme disease	0	0	0	0.00	0.00	0.00
Malaria	0	0	1	0.00	0.00	0.32
Measles	0	0	0	0.00	0.00	0.00
Meningitis, viral	5	18	18	1.28	3.77	5.72
Meningococcal infections	2	5	1	0.51	1.05	0.32
Mumps	4	2	5	1.02	0.42	1.59
Pertussis	2	2	0	0.51	0.42	0.00
Psittacosis	0	0	0	0.00	0.00	0.00
Relapsing fever	0	0	0	0.00	0.00	0.00
Rheumatic fever, acute	0	0	0	0.00	0.00	0.00
Rubella	0	0	0	0.00	0.00	0.00
Salmonellosis	52	69	38	13.28	14.45	12.07
Shigellosis	15	48	16	3.83	10.06	5.08
Strongyloidiasis	0	0	0	0.00	0.00	0.00
Syphilis (early latent)	6	11	15	1.50	2.30	4.80
Syphilis (prim. & secon.)	0	3	1	0.00	0.60	0.30
Syphilis (congenital)	1	2	2	N/A	N/A	N/A
Tetanus	0	1	0	0.00	0.21	0.00
Trichinosis	0	0	0	0.00	0.00	0.00
Tuberculosis	86	59	25	21.96	12.36	7.94
Tularemia	0	0	0	0.00	0.00	0.00
Typhoid fever, case	0	0	2	0.00	0.00	0.64
Typhoid fever, carrier	0	0	0	0.00	0.00	0.00
Typhus fever	2	0	3	0.51	0.00	0.95
Vibrio	2	1	2	0.51	0.21	0.64

^a Rates for perinatal listeriosis were calculated as cases per 100,000 women aged 15 to 44 years.

**Table O-4. Selected Notifiable Diseases
Pomona Health District
Los Angeles County, 1997**

Disease	Frequency	Rate (Cases per 100,000)
	Pomona	Pomona
AIDS	33	5.99
Amebiasis	5	0.91
Botulism	1	0.18
Brucellosis	0	0.00
Campylobacteriosis	68	12.35
Chlamydia	689	152.60
Coccidioidomycosis	3	0.54
Cryptosporidiosis	2	0.36
Cysticercosis	0	0.00
Dengue	1	0.18
E. coli O157:H7	1	0.18
Encephalitis	3	0.54
Giardiasis	21	3.81
Gonorrhea	118	25.40
<i>Haemophilus influenzae</i> type b	0	0.00
Hansen's disease (Leprosy)	0	0.00
Hepatitis A	41	7.45
Hepatitis B	7	1.27
Hepatitis C	0	0.00
Hepatitis unspecified	0	0.00
Kawasaki syndrome	1	0.18
Legionellosis	1	0.18
Listeriosis, nonperinatal	1	0.18
Listeriosis, perinatal ^a	0	0.00
Lyme disease	0	0.00
Malaria	2	0.36
Measles	0	0.00
Meningitis, viral	14	2.54
Meningococcal infections	3	0.54
Mumps	3	0.54
Pertussis	2	0.36
Psittacosis	0	0.00
Relapsing fever	0	0.00
Rheumatic fever, acute	0	0.00
Rubella	0	0.00
Salmonellosis	74	13.44
Shigellosis	23	4.18
Strongyloidiasis	0	0.00
Syphilis (early latent)	13	2.20
Syphilis (prim. & secon.)	2	0.40
Syphilis (congenital)	0	0.00
Tetanus	0	0.00
Trichinosis	0	0.00
Tuberculosis	61	11.08
Tularemia	0	0.00
Typhoid fever, case	1	0.18
Typhoid fever, carrier	0	0.00
Typhus fever	0	0.00
Vibrio	1	0.18

^aRates for perinatal listeriosis were calculated as cases per 100,000 women aged 15 to 44 years.

**Table O-5. Selected Notifiable Diseases
Bellflower, East LA and Whittier Health Districts
Los Angeles County, 1997**

Disease	Frequency			Rate (Cases per 100,000)		
	Bellflower	East LA	Whittier	Bellflower	East LA	Whittier
AIDS	18	34	25	4.98	15.18	7.62
Amebiasis	7	3	1	1.94	1.34	0.30
Botulism	0	0	0	0.00	0.00	0.00
Brucellosis	0	0	0	0.00	0.00	0.00
Campylobacteriosis	39	48	51	10.80	21.43	15.54
Chlamydia	542	515	492	183.00	280.40	184.20
Coccidioidomycosis	1	1	1	0.28	0.45	0.30
Cryptosporidiosis	0	1	0	0.00	0.45	0.00
Cysticercosis	2	3	1	0.55	1.34	0.30
Dengue	0	0	0	0.00	0.00	0.00
E. coli O157:H7	0	0	0	0.00	0.00	0.00
Encephalitis	3	1	1	0.83	0.45	0.30
Giardiasis	26	18	13	7.20	8.04	3.96
Gonorrhea	116	51	62	38.20	27.10	22.60
<i>Haemophilus influenzae</i> type b	1	0	0	0.28	0.00	0.00
Hansen's disease (Leprosy)	0	2	1	0.00	0.89	0.30
Hepatitis A	48	44	49	13.29	19.65	14.93
Hepatitis B	6	1	2	1.66	0.45	0.61
Hepatitis C	1	2	0	0.28	0.89	0.00
Hepatitis unspecified	3	0	0	0.83	0.00	0.00
Kawasaki syndrome	0	1	1	0.00	0.45	0.30
Legionellosis	2	0	0	0.55	0.00	0.00
Listeriosis, nonperinatal	0	0	2	0.00	0.00	0.61
Listeriosis, perinatal ^a	1	0	0	1.24	0.00	0.00
Lyme disease	0	0	0	0.00	0.00	0.00
Malaria	3	0	0	0.83	0.00	0.00
Measles	0	0	0	0.00	0.00	0.00
Meningitis, viral	21	4	9	5.81	1.79	2.74
Meningococcal infections	6	1	3	1.66	0.45	0.91
Mumps	1	1	0	0.28	0.45	0.00
Pertussis	0	1	0	0.00	0.45	0.00
Psittacosis	0	0	0	0.00	0.00	0.00
Relapsing fever	0	0	0	0.00	0.00	0.00
Rheumatic fever, acute	0	0	0	0.00	0.00	0.00
Rubella	4	0	0	1.11	0.00	0.00
Salmonellosis	65	41	70	17.99	18.31	21.33
Shigellosis	24	22	41	6.64	9.82	12.49
Strongyloidiasis	0	0	0	0.00	0.00	0.00
Syphilis (early latent)	8	8	10	2.20	3.60	3.40
Syphilis (prim. & secon.)	0	0	2	0.00	0.00	0.60
Syphilis (congenital)	0	0	0	0.00	0.00	0.00
Tetanus	0	0	0	0.00	0.00	0.00
Trichinosis	0	0	0	0.00	0.00	0.00
Tuberculosis	45	32	59	12.46	14.29	17.98
Tularemia	0	0	0	0.00	0.00	0.00
Typhoid fever, case	1	2	0	0.28	0.89	0.00
Typhoid fever, carrier	0	0	0	0.00	0.00	0.00
Typhus fever	1	0	0	0.28	0.00	0.00
Vibrio	1	0	1	0.28	0.00	0.30

^aRates for perinatal listeriosis were calculated as cases per 100,000 women aged 15 to 44 years.

**Table O-6. Selected Notifiable Diseases
Central and Northeast Health Districts
Los Angeles County, 1997**

Disease	Frequency		Rate (Cases per 100,000)	
	Central	NE	Central	NE
AIDS	202	56	67.55	16.44
Amebiasis	2	5	0.67	1.47
Botulism	0	0	0.00	0.00
Brucellosis	0	1	0.00	0.29
Campylobacteriosis	57	74	19.06	21.72
Chlamydia	934	816	380.50	291.80
Coccidioidomycosis	1	0	0.33	0.00
Cryptosporidiosis	7	1	2.34	0.29
Cysticercosis	2	3	0.67	0.88
Dengue	0	0	0.00	0.00
E. coli O157:H7	0	0	0.00	0.00
Encephalitis	0	0	0.00	0.00
Giardiasis	28	20	9.36	5.87
Gonorrhea	297	108	118.10	37.60
<i>Haemophilus influenzae</i> type b	1	0	0.33	0.00
Hansen's disease (Leprosy)	3	0	1.00	0.00
Hepatitis A	66	76	22.07	22.31
Hepatitis B	3	9	1.00	2.64
Hepatitis C	0	2	0.00	0.59
Hepatitis unspecified	0	0	0.00	0.00
Kawasaki syndrome	0	1	0.00	0.29
Legionellosis	1	0	0.33	0.00
Listeriosis, nonperinatal	0	0	0.00	0.00
Listeriosis, perinatal ^a	0	0	0.00	0.00
Lyme disease	0	0	0.00	0.00
Malaria	6	1	2.01	0.29
Measles	0	0	0.00	0.00
Meningitis, viral	2	0	0.67	0.00
Meningococcal infections	1	3	0.33	0.88
Mumps	0	3	0.00	0.88
Pertussis	2	2	0.67	0.59
Psittacosis	0	0	0.00	0.00
Relapsing fever	0	0	0.00	0.00
Rheumatic fever, acute	0	0	0.00	0.00
Rubella	0	0	0.00	0.00
Salmonellosis	58	61	19.39	17.91
Shigellosis	42	70	14.04	20.55
Strongyloidiasis	0	0	0.00	0.00
Syphilis (early latent)	68	12	22.90	3.50
Syphilis (prim. & secon.)	12	0	4.00	0.00
Syphilis (congenital)	12	3	N/A	N/A
Tetanus	0	0	0.00	0.00
Trichinosis	0	0	0.00	0.00
Tuberculosis	155	55	51.83	16.14
Tularemia	0	0	0.00	0.00
Typhoid fever, case	5	3	1.67	0.88
Typhoid fever, carrier	0	0	0.00	0.00
Typhus fever	0	5	0.00	1.47
Vibrio	1	0	0.33	0.00

^aRates for perinatal listeriosis were calculated as cases per 100,000 women aged 15 to 44 years.

**Table O-7. Selected Notifiable Diseases
Hollywood-Wilshire and West Health Districts
Los Angeles County, 1997**

Disease	Frequency		Rate (Cases per 100,000)	
	Hol-Wil	West	Hol-Wil	West
AIDS	321	71	65.21	11.36
Amebiasis	15	13	3.05	2.08
Botulism	0	0	0.00	0.00
Brucellosis	0	0	0.00	0.00
Campylobacteriosis	66	134	13.41	21.43
Chlamydia	1,159	682	286.80	132.90
Coccidioidomycosis	2	4	0.41	0.64
Cryptosporidiosis	20	8	4.06	1.28
Cysticercosis	1	0	0.20	0.00
Dengue	0	0	0.00	0.00
E. coli O157:H7	0	1	0.00	0.16
Encephalitis	1	1	0.20	0.16
Giardiasis	49	49	9.95	7.84
Gonorrhea	537	200	129.60	38.10
<i>Haemophilus influenzae</i> type b	0	1	0.00	0.16
Hansen's disease (Leprosy)	0	0	0.00	0.00
Hepatitis A	203	66	41.24	10.56
Hepatitis B	17	2	3.45	0.32
Hepatitis C	2	0	0.41	0.00
Hepatitis unspecified	0	0	0.00	0.00
Kawasaki syndrome	0	0	0.00	0.00
Legionellosis	1	10	0.20	1.60
Listeriosis, nonperinatal	3	1	0.61	0.16
Listeriosis, perinatal ^a	2	3	1.87	2.05
Lyme disease	0	1	0.00	0.17
Malaria	4	9	0.81	1.44
Measles	1	0	0.20	0.00
Meningitis, viral	5	1	1.02	0.16
Meningococcal infections	4	4	0.81	0.64
Mumps	1	0	0.20	0.00
Pertussis	4	2	0.81	0.32
Psittacosis	0	0	0.00	0.00
Relapsing fever	0	0	0.00	0.00
Rheumatic fever, acute	0	0	0.00	0.00
Rubella	0	0	0.00	0.00
Salmonellosis	101	107	20.52	17.11
Shigellosis	65	49	13.20	7.84
Strongyloidiasis	2	0	0.41	0.00
Syphilis (early latent)	54	11	10.90	1.60
Syphilis (prim. & secon.)	9	4	1.80	0.60
Syphilis (congenital)	2	2	N/A	N/A
Tetanus	0	0	0.00	0.00
Trichinosis	0	0	0.00	0.00
Tuberculosis	116	53	23.56	8.48
Tularemia	0	1	0.00	0.16
Typhoid fever, case	3	1	0.61	0.16
Typhoid fever, carrier	0	0	0.00	0.00
Typhus fever	0	0	0.00	0.00
Vibrio	2	7	0.41	1.12

^aRates for perinatal listeriosis were calculated as cases per 100,000 women aged 15 to 44 years.

**Table O-8. Selected Notifiable Diseases
Compton, San Antonio and South Health Districts
Los Angeles County, 1997**

Disease	Frequency			Rate (Cases per 100,000)		
	Comp	S Antonio	South	Comp	S Antonio	South
AIDS	27	52	35	9.69	12.19	20.92
Amebiasis	7	10	1	2.51	2.34	0.60
Botulism	0	0	0	0.00	0.00	0.00
Brucellosis	0	2	0	0.00	0.47	0.00
Campylobacteriosis	30	51	18	10.77	11.96	10.76
Chlamydia	1,188	982	994	519.80	280.60	723.80
Coccidioidomycosis	0	0	0	0.00	0.00	0.00
Cryptosporidiosis	0	1	0	0.00	0.23	0.00
Cysticercosis	1	3	0	0.36	0.70	0.00
Dengue	0	0	0	0.00	0.00	0.00
E. coli O157:H7	0	0	0	0.00	0.00	0.00
Encephalitis	1	3	2	0.36	0.70	1.20
Giardiasis	36	30	14	12.92	7.03	8.37
Gonorrhea	358	94	446	152.90	26.30	316.80
<i>Haemophilus influenzae</i> type b	1	0	0	0.35	0.00	0.00
Hansen's disease (Leprosy)	1	1	0	0.36	0.23	0.00
Hepatitis A	53	88	33	19.02	20.63	19.72
Hepatitis B	6	1	5	2.15	0.23	2.99
Hepatitis C	0	1	4	0.00	0.23	2.39
Hepatitis unspecified	0	1	0	0.00	0.23	0.00
Kawasaki syndrome	1	0	0	0.36	0.00	0.00
Legionellosis	1	2	0	0.36	0.47	0.00
Listeriosis, nonperinatal	1	0	0	0.36	0.00	0.00
Listeriosis, perinatal ^a	0	0	1	0.00	0.00	2.73
Lyme disease	0	0	0	0.00	0.00	0.00
Malaria	1	1	0	0.36	0.23	0.00
Measles	1	0	1	0.36	0.00	0.60
Meningitis, viral	9	22	6	3.23	5.16	3.59
Meningococcal infections	5	1	2	1.79	0.23	1.20
Mumps	1	0	0	0.36	0.00	0.00
Pertussis	1	1	1	0.36	0.23	0.60
Psittacosis	0	0	0	0.00	0.00	0.00
Relapsing fever	0	0	0	0.00	0.00	0.00
Rheumatic fever, acute	0	0	1	0.00	0.00	0.60
Rubella	0	0	0	0.00	0.00	0.00
Salmonellosis	46	64	41	16.51	15.01	24.50
Shigellosis	33	60	20	11.85	14.07	11.95
Strongyloidiasis	0	0	0	0.00	0.00	0.00
Syphilis (early latent)	49	27	84	18.40	6.60	48.80
Syphilis (prim. & secon.)	6	3	22	2.20	0.70	13.10
Syphilis (congenital)	6	1	8	N/A	N/A	N/A
Tetanus	0	0	0	0.00	0.00	0.00
Trichinosis	1	1	0	0.36	0.23	0.00
Tuberculosis	42	57	46	15.08	13.36	27.49
Tularemia	0	0	0	0.00	0.00	0.00
Typhoid fever, case	0	3	0	0.00	0.70	0.00
Typhoid fever, carrier	0	0	0	0.00	0.00	0.00
Typhus fever	0	0	0	0.00	0.00	0.00
Vibrio	0	1	1	0.00	0.23	0.60

^aRates for perinatal listeriosis were calculated as cases per 100,000 women aged 15 to 44 years.

**Table O-9. Selected Notifiable Diseases
Southeast, Southwest, and Inglewood Health Districts
Los Angeles County, 1997**

Disease	Frequency			Rate (Cases per 100,000)		
	Ing	SE	SW	Ing	SE	SW
AIDS	59	34	117	14.32	22.90	32.30
Amebiasis	4	0	4	0.97	0.00	1.10
Botulism	0	0	0	0.00	0.00	0.00
Brucellosis	0	0	0	0.00	0.00	0.00
Campylobacteriosis	48	30	41	11.65	20.20	11.32
Chlamydia	1,553	634	1,836	459.50	520.60	617.90
Coccidioidomycosis	1	0	0	0.24	0.00	0.00
Cryptosporidiosis	3	2	2	0.73	1.35	0.55
Cysticercosis	0	3	1	0.00	2.02	0.28
Dengue	0	0	0	0.00	0.00	0.00
E. coli O157:H7	0	0	0	0.00	0.00	0.00
Encephalitis	0	1	0	0.00	0.67	0.00
Giardiasis	36	21	13	8.74	14.14	3.59
Gonorrhea	605	192	770	174.50	153.50	252.60
<i>Haemophilus influenzae</i> type b	3	0	1	0.73	0.00	0.28
Hansen's disease (Leprosy)	1	0	0	0.24	0.00	0.00
Hepatitis A	56	43	46	13.59	28.96	12.70
Hepatitis B	9	2	2	2.18	1.35	0.55
Hepatitis C	0	0	0	0.00	0.00	0.00
Hepatitis unspecified	1	0	1	0.24	0.00	0.28
Kawasaki syndrome	0	0	1	0.00	0.00	0.28
Legionellosis	4	0	1	0.97	0.00	0.28
Listeriosis, nonperinatal	0	0	1	0.00	0.00	0.28
Listeriosis, perinatal ^a	0	0	0	0.00	0.00	0.00
Lyme disease	0	0	0	0.00	0.00	0.00
Malaria	3	0	1	0.73	0.00	0.28
Measles	0	0	1	0.00	0.00	0.28
Meningitis, viral	7	6	11	1.70	4.04	3.04
Meningococcal infections	5	2	3	1.21	1.35	0.83
Mumps	3	1	0	0.73	0.67	0.00
Pertussis	1	1	2	0.24	0.67	0.55
Psittacosis	0	0	0	0.00	0.00	0.00
Relapsing fever	0	0	0	0.00	0.00	0.00
Rheumatic fever, acute	0	0	0	0.00	0.00	0.00
Rubella	0	0	0	0.00	0.00	0.00
Salmonellosis	79	29	83	19.18	19.53	22.92
Shigellosis	29	28	34	7.04	18.86	9.39
Strongyloidiasis	1	1	0	0.24	0.67	0.00
Syphilis (early latent)	81	40	87	20.10	27.10	24.20
Syphilis (prim. & secon.)	14	2	15	3.40	1.30	4.10
Syphilis (congenital)	9	10	8	N/A	N/A	N/A
Tetanus	0	0	1	0.00	0.00	0.28
Trichinosis	0	0	0	0.00	0.00	0.00
Tuberculosis	60	25	72	14.57	16.84	19.88
Tularemia	0	0	0	0.00	0.00	0.00
Typhoid fever, case	1	0	0	0.24	0.00	0.00
Typhoid fever, carrier	0	0	0	0.00	0.00	0.00
Typhus fever	0	0	0	0.00	0.00	0.00
Vibrio	0	0	0	0.00	0.00	0.00

^aRates for perinatal listeriosis were calculated as cases per 100,000 women aged 15 to 44 years.

**Table O-10. Selected Notifiable Diseases
Harbor and Torrance Health Districts
Los Angeles County, 1997**

Disease	Frequency		Rate (Cases per 100,000)	
	Harbor	Torrance	Harbor	Torrance
AIDS	11	16	5.45	3.50
Amebiasis	4	9	1.98	1.97
Botulism	0	0	0.00	0.00
Brucellosis	0	0	0.00	0.00
Campylobacteriosis	59	126	29.23	27.59
Chlamydia	400	591	241.30	157.70
Coccidioidomycosis	0	1	0.00	0.22
Cryptosporidiosis	1	3	0.50	0.66
Cysticercosis	1	0	0.50	0.00
Dengue	0	0	0.00	0.00
E. coli O157:H7	0	4	0.00	0.88
Encephalitis	1	1	0.50	0.22
Giardiasis	29	75	14.37	16.43
Gonorrhea	80	140	47.10	36.40
<i>Haemophilus influenzae</i> type b	0	0	0.00	0.00
Hansen's disease (Leprosy)	0	3	0.00	0.66
Hepatitis A	51	47	25.27	10.29
Hepatitis B	1	6	0.50	1.31
Hepatitis C	1	1	0.50	0.22
Hepatitis unspecified	0	1	0.00	0.22
Kawasaki syndrome	0	1	0.00	0.22
Legionellosis	1	2	0.50	0.44
Listeriosis, nonperinatal	1	0	0.50	0.00
Listeriosis, perinatal ^a	0	0	0.00	0.00
Lyme disease	1	1	0.50	0.22
Malaria	0	4	0.00	0.88
Measles	0	0	0.00	0.00
Meningitis, viral	9	6	4.46	1.31
Meningococcal infections	0	3	0.00	0.66
Mumps	1	5	0.50	1.10
Pertussis	0	1	0.00	0.22
Psittacosis	1	0	0.50	0.00
Relapsing fever	0	0	0.00	0.00
Rheumatic fever, acute	0	0	0.00	0.00
Rubella	0	0	0.00	0.00
Salmonellosis	29	121	14.37	26.50
Shigellosis	23	31	11.40	6.79
Strongyloidiasis	0	0	0.00	0.00
Syphilis (early latent)	4	3	2.00	0.70
Syphilis (prim. & secon.)	0	0	0.00	0.00
Syphilis (congenital)	0	0	0.00	0.00
Tetanus	0	0	0.00	0.00
Trichinosis	0	0	0.00	0.00
Tuberculosis	11	45	5.45	9.86
Tularemia	0	0	0.00	0.00
Typhoid fever, case	1	1	0.50	0.22
Typhoid fever, carrier	0	0	0.00	0.00
Typhus fever	0	0	0.00	0.00
Vibrio	0	1	0.00	0.22

^aRates for perinatal listeriosis were calculated as cases per 100,000 women aged 15 to 44 years.