| Due Due sue en en el le elle | | | | | | | | | | | | | | | | |
|---|------------------|-----|------------------|-----|------------------|-----|------------------|-----|----------------|--------|------------------|-----|------------------|-----|------------------|--------|
| Pre-Pregnancy Health | | | | | | | | | | | | | | | | |
| by Service Planning Area | | | | | | | | | | | | | | | | |
| by service riaming / irea | Antelope Valley | | San Fernando | | San Gabriel | | Metro | | West | | South | | East | | South Bay | |
| | SPA 1 | | SPA 2 | | SPA 3 | | SPA 4 | | SPA 5 | | SPA 6 | | SPA 7 | | SPA 8 | |
| | % 95% CI | N | % 95% C | N | % 95% CI | N | % 95% CI | N | % 95% CI | N |
| Access to Health Care Before Pregnancy | | | | | | | | | | | | | | | | |
| Insured | 90.3 (80.4-100) | 154 | 87.2 (80.7-93.6) | 423 | 87.0 (80.5-93.4) | 431 | 91.3 (84.5-98.2) | 224 | 100.0 (100-10 | 0) 108 | 88.0 (82.2-93.8) | 476 | 84.6 (76.6-92.5) | 337 | 83.2 (72.5-93.8 | 319 |
| On Medi-Cal | 52.8 (35.5-70.2 | 81 | 38.3 (27.7-48.9) | 153 | 37.5 (27.7-47.3) | 161 | 38.7 (24.8-52.5) | 85 | 12.1* (0-24.8 |) 13 | 70.6 (60.4-80.8) | 330 | 51.2 (38.5-63.9) | 170 | 36.0 (22.7-49.3 |) 113 |
| Preconception Health & Well-being | | | | | | | | | | | | | | | | |
| No preconception health counseling ¹ | 60.9 (44.1-77.6 | 101 | 56.8 (47.1-66.4) | 276 | 60.2 (50.7-69.6) | 292 | 47.1 (33.9-60.3) | 120 | 41.6 (22.3-60 | 9) 45 | 63.6 (53.8-73.4) | 351 | 67.9 (57.1-78.8) | 273 | 47.9 (35-60.8) | 180 |
| Not taking a multivitamin | 62.1 (46.1-78.1 | 103 | 43.3 (33.7-52.8) | 212 | 42.8 (33.4-52.2) | 213 | 30.8 (18.9-42.7) | 78 | 31.5* (13.2-49 | 9) 34 | 47.3 (37.3-57.2) | 267 | 54.2 (42.8-65.5) | 217 | 42.9 (30.2-55.6 |) 170 |
| Lack of folic acid knowledge | 70.2 (54.3-86.1 | 119 | 63.5 (54-73) | 299 | 58.1 (48.6-67.5) | 288 | 55.9 (42.4-69.5) | 137 | 57.2 (38.5-75 | 8) 60 | 75.6 (67.2-84) | 412 | 59.7 (48.2-71.1) | 229 | 65.4 (52.5-78.2 | 251 |
| Had enough money for food | 91.8 (82.2-100) | 156 | 98.2 (95.6-100) | 481 | 97.1 (94.3-100) | 485 | 96.8 (92.3-100) | 247 | 97.2 (91.8-10 | 0) 105 | 94.2 (89.4-99.1) | 539 | 98.4 (95.2-100) | 397 | 94.9 (89.8-100) |) 377 |
| Did not smoke in the 6 months before pregnancy | 85.1 (73.5-96.6 | 142 | 94.7 (90.3-99) | 464 | 96.7 (93.7-99.6) | 478 | 91.8 (84.6-99) | 234 | 93.8 (86.3-10 | 0) 93 | 87.3 (80.2-94.4) | 487 | 94.6 (89.8-99.4) | 382 | 88.7 (81.8-95.6 | 340 |
| Family Planning Practices | | | | | | | | | | | | | | | | |
| Unwanted pregnancy/mistimed by mother ² | 48.5 (32.1-65) | 79 | 31.7 (22.7-40.7) | 149 | 35.4 (26-44.8) | 170 | 20.6* (10-31.3) | 51 | 21.3* (2.2-40. | 5) 22 | 37.7 (27.9-47.4) | 202 | 32.7 (22.1-43.2) | 129 | 29.3 (16.8-41.8 | 3) 111 |
| Unwanted pregnancy by partner ³ | 26.0* (11.9-40.1 | 44 | 24.2 (16-32.4) | 113 | 17.2 (10.1-24.3) | 84 | 17.0* (7.2-26.8) | 43 | 24.2* (4.5-43. | 3) 24 | 22.6 (14.3-31) | 125 | 23.1 (13.6-32.6) | 92 | 15.7* (6.5-24.9) | 59 |
| Using birth control before getting pregnant⁴ | 28.7* (14.4-43) | 49 | 23.3 (15.2-31.5) | 113 | 31.0 (22.1-39.9) | 152 | 26.1 (14.6-37.6) | 65 | 9.4* (0.5-18. | 3) 10 | 24.6 (16.2-33) | 136 | 33.5 (22.6-44.5) | 131 | 28.4 (16.1-40.7 |) 112 |
| Preconception Health Conditions | | | | | | | | | | | | | | | | |
| Overweight/Obese | 73.9 (59.8-88.1 | 120 | 49.0 (39-59) | 226 | 49.9 (40.2-59.7) | 233 | 48.7 (35.3-62.1) | 123 | 25.0* (8.6-41. | 4) 25 | 71.5 (62.2-80.8) | 385 | 80.7 (71.8-89.5) | 288 | 55.6 (42.2-69.1 | .) 206 |
| Did not have asthma | 81.8 (69.3-94.3 | 139 | 90.5 (84.5-96.6) | 444 | 94.3 (90.2-98.3) | 470 | 92.5 (84.6-100) | 236 | 93.7 (85.1-10 | 0) 101 | 84.4 (77-91.9) | 483 | 97.7 (95-100) | 395 | 92.9 (87.1-98.6 | 369 |
| Did not have high blood pressure | 82.9 (70.9-95) | 141 | 82.9 (75.6-90.3) | 406 | 92.8 (88.8-96.8) | 463 | 85.4 (75.1-95.7) | 218 | 85.6 (70.9-10 | 0) 92 | 87.5 (80.9-94.2) | 501 | 91.1 (82.6-99.5) | 368 | 86.8 (76.8-96.7 |) 344 |
| Did not have high blood sugar | 91.0 (80.8-100) | 155 | 94.0 (89.6-98.5) | 461 | 93.1 (88.4-97.7) | 464 | 100.0 (100-100) | 255 | 90.74 (80.3-10 | 0) 98 | 88.1 (81.5-94.8) | 504 | 94.6 (90.3-98.9) | 382 | 93.9 (88.8-99.1 | .) 373 |
| Did not have anemia | 95.3 (88.9-100) | 162 | 98.6 (96.6-100) | 483 | 94.3 (90-98.7) | 471 | 98.2 (94.6-100) | 250 | 100.0 (100-10 | 0) 108 | 95.5 (91.8-99.3) | 546 | 94.4 (89.2-99.5) | 381 | 95.2 (90.7-99.8 | 378 |
| Did not have periodontal disease | 100.0 (100-100) | 170 | 100.0 (100-100) | 490 | 99.2 (97.7-100) | 495 | 100.0 (100-100) | 255 | 100.0 (100-10 |) 108 | 96.6 (92.6-100) | 553 | 100.0 (100-100) | 404 | 99.0 (97-100) | 393 |
| Did not report depression | 90.3 (80.4-100) | 154 | 95.7 (92.2-99.3) | 469 | 96.1 (93.2-99.1) | 480 | 98.2 (95.7-100) | 250 | 99.7 (99.1-10 | 0) 108 | 92.9 (87.7-98.1) | 531 | 90.0 (81.6-98.3) | 363 | 100.0 (100-100) | 397 |
| Did not report anxiety | 82.6 (70.3-94.8 | 140 | 97.2 (94.3-100) | 476 | 95.6 (92-99.2) | 477 | 95.6 (90.1-100) | 244 | 95.1 (87.7-10 | 0) 103 | 94.0 (89.2-98.8) | 538 | 100.0 (100-100) | 404 | 86.0 (76.8-95.2 | 341 |
| Obstetric History | | | | | | | | | | | | | | | | |
| Did not have a preterm birth previously | 85.4 (73.4-97.5 | 126 | 87.3 (80.1-94.5) | 366 | 82.9 (74.8-91) | 329 | 88.5 (78.8-98.3) | 196 | 90.8 (78.5-10 | 0) 83 | 73.4 (63.2-83.5) | 341 | 78.6 (68.6-88.5) | 266 | 93.3 (86.4-100) |) 330 |
| Did not have a low birth weight baby previously | 81.7 (68.5-95) | 121 | 94.9 (90.5-99.3) | 372 | 85.8 (78.2-93.5) | 331 | 90.0 (82-98) | 197 | 95.6 (87.3-10 | 0) 88 | 78.4 (68.7-88.1) | 341 | 85.5 (77.3-93.6) | 294 | 92.5 (85.2-99.8 | 322 |
| Did not experience a stillbirth (pregnancy that did not result in a live birth) | 81.6 (67.7-95.6 | 115 | 89.4 (82.4-96.5) | 381 | 95.1 (91-99.2) | 368 | 88.6 (80.2-96.9) | 198 | 100.0 (100-10 | 0) 84 | 84.5 (76.5-92.5) | 383 | 94.8 (89.5-100) | 317 | 91.3 (83.3-99.2 | 2) 315 |
| Did not have a miscarriage previously | 61.9 (45.1-78.6 | 103 | 82.7 (74.9-90.5) | 352 | 72.2 (63.1-81.4) | 308 | 72.7 (60.1-85.3) | 161 | 75.9 (59.2-92 | 5) 70 | 66.2 (55.7-76.6) | 316 | 67.4 (56.7-78.1) | 241 | 74.3 (63.1-85.5 |) 280 |
| Did not have an abortion previously | 80.2 (67.3-93) | 116 | 83.8 (76.1-91.5) | 353 | 87.0 (79.7-94.4) | 342 | 81.8 (71-92.6) | 176 | 77.0 (60.8-93 | 1) 73 | 71.4 (60.5-82.3) | 322 | 96.1 (92.3-100) | 323 | 84.3 (75.1-93.4 |) 291 |
| Did not experience infant death previously | 90.2 (78.6-100) | 124 | 98.0 (95.3-100) | 407 | 97.1 (93.9-100) | 388 | 96.3 (91.2-100) | 214 | 100.0 (100-10 | 0) 88 | 91.4 (84.5-98.4) | 399 | 96.5 (92.2-100) | 330 | 94.8 (87-100) | 327 |
| | | | | | | _ | | | | | | | | | | |

^{*}Signifies that the estimate is statistically unstable (relative standard error > 25%) and therefore may not be appropriate to use for planning or policy purposes.

^{&#}x27;Question asked: During the six months before you got pregnant with your new baby, did you talk to a doctor, nurse, or other health care worker about how to prepare for a healthy baby and pregnancy?

²Question asked: Thinking back to just before you got pregnant with your last baby, how did you feel about becoming pregnant? Respondents who answered "I wanted to be pregnant later" or "I didn't want to be pregnant then or any time in the future" were coded as "Unwanted or mistimed."

³Question asked: Just before you got pregnant with your new baby, how did your husband or partner feel about you becoming pregnant? Respondents who answered, "I don't know" or "I didn't have a husband or partner" were coded missing for this field.

^{*}Question asked: Before you got pregnant, were you doing anything to keep from getting pregnant? Respondents who answered "Yes, all the time" or "Yes, sometimes" were coded as "Using birth control before getting pregnant."