

## 2010-2012 LA HOPE Surveys

Domain/Subdomain		<i>Nutrition &amp; Physical Activity / Obesity and Overweight</i>			<i>Nutrition &amp; Physical Activity / Folic Acid Supplementation</i>			<i>Nutrition &amp; Physical Activity / Exercise/Physical Activity</i>		
Indicator Description		Percentage of women having an infant/fetal death who are overweight* or obese** based on body mass index (BMI) *Overweight: percentage of women with a pre-pregnancy BMI $\geq$ 25 kg/m <sup>2</sup> but < 30 kg/m <sup>2</sup> **Obesity: percentage of women with a pre-pregnancy BMI $\geq$ 30 kg/m <sup>2</sup>			Percentage of women having an infant/fetal death who did not take a multivitamin, prenatal vitamin, or a folic acid supplement during one month prior to pregnancy.			Percentage of women having an infant/fetal death who did not exercise during pregnancy.		
		%	95% CI		%	95% CI		%	95% CI	
<b>LA County</b>		56.5	52.8	60.2	50.2	46.7	53.8	41.6	38.0	45.3
<b>Race/Ethnicity</b>	<b>White</b>	48.6	40.0	57.2	31.3	23.5	39.2	40.7	32.0	49.3
	<b>Latina</b>	60.9	56.3	65.4	56.7	52.4	61.0	41.0	36.5	45.5
	<b>African-American</b>	65.4	56.2	74.6	53.4	43.9	62.9	37.1	27.7	46.4
	<b>Asian/Pacific Islander</b>	34.0	17.7	50.3	37.2	22.3	52.0	51.6	35.8	67.4
<b>Service Planning Areas</b>	<b>SPA 1 Antelope Valley</b>	70.5	59.7	81.4	54.2	42.7	65.6	43.3	31.6	55.0
	<b>SPA 2 San Fernando</b>	53.4	44.6	62.1	35.5	27.7	43.4	47.0	38.1	55.9
	<b>SPA 3 San Gabriel</b>	47.0	37.2	56.7	48.5	39.1	57.8	43.3	33.9	52.7
	<b>SPA 4 Metro</b>	57.8	45.8	69.7	51.2	40.1	62.4	30.9	20.6	41.1
	<b>SPA 5 West</b>	22.4*	10.3	34.6	26.8*	13.5	40.2	40.6	24.4	56.8
	<b>SPA 6 South</b>	65.9	58.2	73.6	63.8	56.3	71.2	31.7	24.3	39.2
	<b>SPA 7 East</b>	63.2	53.7	72.7	56.9	47.8	66.1	48.9	39.2	58.7
	<b>SPA 8 South Bay</b>	53.7	42.7	64.7	49.6	39.0	60.1	46.4	35.7	57.1

\*The estimate is statistically unstable (relative standard error > 25%) and therefore may not be appropriate to use for planning or policy purposes.

[For more information about LA HOPE project please click here](#)